

Chapter 6

A Tour of the Cell

PowerPoint® Lecture Presentations for

Biology

Eighth Edition

Neil Campbell and Jane Reece

Lectures by Chris Romero, updated by Erin Barley with contributions from Joan Sharp

Overview: The Fundamental Units of Life

- All organisms are made of cells
- The cell is the simplest collection of matter that can live
- Cell structure is correlated to cellular function
- All cells are related by their descent from earlier cells

Concept 6.1: To study cells, biologists use microscopes and the tools of biochemistry


- Though usually too small to be seen by the unaided eye, cells can be complex

Microscopy

- Scientists use microscopes to visualize cells too small to see with the naked eye
- In a **light microscope (LM)**, visible light passes through a specimen and then through glass lenses, which magnify the image

-
- The quality of an image depends on
 - *Magnification*, the ratio of an object's image size to its real size
 - *Resolution*, the measure of the clarity of the image, or the minimum distance of two distinguishable points
 - *Contrast*, visible differences in parts of the sample

Fig. 6-2



-
- LMs can magnify effectively to about 1,000 times the size of the actual specimen
 - Various techniques enhance contrast and enable cell components to be stained or labeled
 - Most subcellular structures, including **organelles** (membrane-enclosed compartments), are too small to be resolved by an LM

Fig. 6-3

TECHNIQUE

RESULTS

(a) Brightfield (unstained specimen)

50 μ m

(b) Brightfield (stained specimen)

(c) Phase-contrast

(d) Differential-interference-contrast (Nomarski)

(e) Fluorescence

50 μ m

(f) Confocal

50 μ m

TECHNIQUE

(a) Brightfield (unstained specimen)

RESULTS

(b) Brightfield (stained specimen)

50 μm

TECHNIQUE

(c) Phase-contrast

RESULTS

(d) Differential-interference-contrast (Nomarski)

TECHNIQUE

(e) Fluorescence

RESULTS





Fig. 6-3f

TECHNIQUE

RESULTS

(f)
Confoca
I



50 μm


-
- Two basic types of **electron microscopes (EMs)** are used to study subcellular structures
 - **Scanning electron microscopes (SEMs)** focus a beam of electrons onto the surface of a specimen, providing images that look 3-D
 - **Transmission electron microscopes (TEMs)** focus a beam of electrons through a specimen
 - TEMs are used mainly to study the internal structure of cells

TECHNIQUE


RESULTS



**(a) Scanning electron
microscopy (SEM)**

Cilia


1 μm


**(b) Transmission electron
microscopy (TEM)**

**Longitudinal
section of
cilium**


**Cross section
of cilium**

1 μm


Cell Fractionation

- **Cell fractionation** takes cells apart and separates the major organelles from one another
- Ultracentrifuges fractionate cells into their component parts
- Cell fractionation enables scientists to determine the functions of organelles
- Biochemistry and cytology help correlate cell function with structure

Fig. 6-5

TECHNIQUE




Fig. 6-5a

TECHNIQUE

Homogenization

**Tissue
cells**



Homogenate



Differential centrifugation

Fig. 6-5b

TECHNIQUE (cont.)

1,000 g
(1,000 times the
force of gravity)
10 min

Supernatant poured
into next tube

20,000 g
20 min

|
Pellet rich in
nuclei and
cellular debris

80,000 g
60 min

|
Pellet rich in
mitochondria
(and
chloro-plasts if
cells
are from a plant)

150,000 g
3 hr

|
Pellet rich in
“microsomes”
(pieces of plasma
membranes and
cells’ internal
membranes)

|
Pellet rich in
ribosomes


Concept 6.2: Eukaryotic cells have internal membranes that compartmentalize their functions

- The basic structural and functional unit of every organism is one of two types of cells: prokaryotic or eukaryotic
- Only organisms of the domains Bacteria and Archaea consist of prokaryotic cells
- Protists, fungi, animals, and plants all consist of eukaryotic cells

Comparing Prokaryotic and Eukaryotic Cells

- Basic features of all cells:
 - Plasma membrane
 - Semifluid substance called **cytosol**
 - Chromosomes (carry genes)
 - Ribosomes (make proteins)

-
- **Prokaryotic cells** are characterized by having
 - No nucleus
 - DNA in an unbound region called the **nucleoid**
 - No membrane-bound organelles
 - **Cytoplasm** bound by the plasma membrane




-
- **Eukaryotic cells** are characterized by having
 - DNA in a nucleus that is bounded by a membranous nuclear envelope
 - Membrane-bound organelles
 - Cytoplasm in the region between the plasma membrane and nucleus
 - Eukaryotic cells are generally much larger than prokaryotic cells

-
- The **plasma membrane** is a selective barrier that allows sufficient passage of oxygen, nutrients, and waste to service the volume of every cell
 - The general structure of a biological membrane is a double layer of phospholipids

Outside of cell

(a) TEM of a plasma membrane



(b) Structure of the plasma membrane

-
- The logistics of carrying out cellular metabolism sets limits on the size of cells
 - The surface area to volume ratio of a cell is critical
 - As the surface area increases by a factor of n^2 , the volume increases by a factor of n^3
 - Small cells have a greater surface area relative to volume

Fig. 6-8

**Surface area increases while
total volume remains constant**

	1	5	1
Total surface area [Sum of the surface areas (height × width) of all boxes sides × number of boxes]	6	150	750
Total volume [height × width × length × number of boxes]	1	125	125
Surface-to-volume (S-to-V) ratio [surface area ÷ volume]	6	1.2	6

A Panoramic View of the Eukaryotic Cell

- A eukaryotic cell has internal membranes that partition the cell into organelles
- Plant and animal cells have most of the same organelles

BioFlix: Tour Of An Animal Cell

BioFlix: Tour Of A Plant Cell

Fig. 6-9a





Fig. 6-9b



Concept 6.3: The eukaryotic cell's genetic instructions are housed in the nucleus and carried out by the ribosomes





- The nucleus contains most of the DNA in a eukaryotic cell
- Ribosomes use the information from the DNA to make proteins

The Nucleus: Information Central

- The **nucleus** contains most of the cell's genes and is usually the most conspicuous organelle
- The **nuclear envelope** encloses the nucleus, separating it from the cytoplasm
- The nuclear membrane is a double membrane; each membrane consists of a lipid bilayer

Fig. 6-10

1 μm



Surface of nuclear envelope

Ribosome

1 μm

0.25 μm

Close-up of nuclear envelope

Pore complexes (TEM)

Nuclear lamina (TEM)

-
- Pores regulate the entry and exit of molecules from the nucleus
 - The shape of the nucleus is maintained by the **nuclear lamina**, which is composed of protein

-
- In the nucleus, DNA and proteins form genetic material called **chromatin**
 - Chromatin condenses to form discrete **chromosomes**
 - The **nucleolus** is located within the nucleus and is the site of ribosomal RNA (rRNA) synthesis

Ribosomes: Protein Factories

- **Ribosomes** are particles made of ribosomal RNA and protein
- Ribosomes carry out protein synthesis in two locations:
 - In the cytosol (free ribosomes)
 - On the outside of the endoplasmic reticulum or the nuclear envelope (bound ribosomes)

— Cytosol

⌋ Endoplasmic reticulum (ER)

Free ribosomes

Bound ribosomes

Large subunit

Small subunit

0.5 μm

TEM showing ER and ribosomes

Diagram of a ribosome


Concept 6.4: The endomembrane system regulates protein traffic and performs metabolic functions in the cell

- Components of the **endomembrane system**:
 - Nuclear envelope
 - Endoplasmic reticulum
 - Golgi apparatus
 - Lysosomes
 - Vacuoles
 - Plasma membrane
- These components are either continuous or connected via transfer by **vesicles**

The Endoplasmic Reticulum: Biosynthetic Factory

- The **endoplasmic reticulum (ER)** accounts for more than half of the total membrane in many eukaryotic cells
- The ER membrane is continuous with the nuclear envelope
- There are two distinct regions of ER:
 - **Smooth ER**, which lacks ribosomes
 - **Rough ER**, with ribosomes studding its surface

Fig. 6-12



Functions of Smooth ER

- The smooth ER
 - Synthesizes lipids
 - Metabolizes carbohydrates
 - Detoxifies poison
 - Stores calcium

Functions of Rough ER

- The rough ER
 - Has bound ribosomes, which secrete **glycoproteins** (proteins covalently bonded to carbohydrates)
 - Distributes **transport vesicles**, proteins surrounded by membranes
 - Is a membrane factory for the cell

The Golgi Apparatus: Shipping and Receiving Center

- The **Golgi apparatus** consists of flattened membranous sacs called cisternae
- Functions of the Golgi apparatus:
 - Modifies products of the ER
 - Manufactures certain macromolecules
 - Sorts and packages materials into transport vesicles

Fig. 6-13

cis face
("receiving" side of
Golgi apparatus)

Cisternae

0.1 μm

trans face
("shipping" side of
Golgi apparatus)

TEM of Golgi apparatus



Lysosomes: Digestive Compartments

- A **lysosome** is a membranous sac of hydrolytic enzymes that can digest macromolecules
- Lysosomal enzymes can hydrolyze proteins, fats, polysaccharides, and nucleic acids

Animation: Lysosome Formation

-
- Some types of cell can engulf another cell by **phagocytosis**; this forms a food vacuole
 - A lysosome fuses with the food vacuole and digests the molecules
 - Lysosomes also use enzymes to recycle the cell's own organelles and macromolecules, a process called autophagy

Fig. 6-14






Fig. 6-14a



(a) Phagocytosis

Fig. 6-14b



(b) Autophagy


Vacuoles: Diverse Maintenance Compartments

- A plant cell or fungal cell may have one or several vacuoles

-
- **Food vacuoles** are formed by phagocytosis
 - **Contractile vacuoles**, found in many freshwater protists, pump excess water out of cells
 - **Central vacuoles**, found in many mature plant cells, hold organic compounds and water

[Video: Paramecium Vacuole](#)

Fig. 6-15



The Endomembrane System: *A Review*

- The endomembrane system is a complex and dynamic player in the cell's compartmental organization

Fig. 6-16-1

Smooth ER

Nucleus

Rough ER

Plasma
membrane

Fig. 6-16-2





Fig. 6-16-3



Concept 6.5: Mitochondria and chloroplasts change energy from one form to another


- **Mitochondria** are the sites of cellular respiration, a metabolic process that generates ATP
- **Chloroplasts**, found in plants and algae, are the sites of photosynthesis
- **Peroxisomes** are oxidative organelles

-
- Mitochondria and chloroplasts
 - Are not part of the endomembrane system
 - Have a double membrane
 - Have proteins made by free ribosomes
 - Contain their own DNA

Mitochondria: Chemical Energy Conversion

- Mitochondria are in nearly all eukaryotic cells
- They have a smooth outer membrane and an inner membrane folded into **cristae**
- The inner membrane creates two compartments: intermembrane space and **mitochondrial matrix**
- Some metabolic steps of cellular respiration are catalyzed in the mitochondrial matrix
- Cristae present a large surface area for enzymes that synthesize ATP

Fig. 6-17




0.1 μm

Chloroplasts: Capture of Light Energy

- The chloroplast is a member of a family of organelles called **plastids**
- Chloroplasts contain the green pigment chlorophyll, as well as enzymes and other molecules that function in photosynthesis
- Chloroplasts are found in leaves and other green organs of plants and in algae

-
- Chloroplast structure includes:
 - **Thylakoids**, membranous sacs, stacked to form a **granum**
 - **Stroma**, the internal fluid


Fig. 6-18



Peroxisomes: Oxidation

- Peroxisomes are specialized metabolic compartments bounded by a single membrane
- Peroxisomes produce hydrogen peroxide and convert it to water
- Oxygen is used to break down different types of molecules

Fig. 6-19




1 μm

A scale bar consisting of a horizontal line with vertical end caps, labeled '1 μm ' below it.

Concept 6.6: The cytoskeleton is a network of fibers that organizes structures and activities in the cell


- The **cytoskeleton** is a network of fibers extending throughout the cytoplasm
- It organizes the cell's structures and activities, anchoring many organelles
- It is composed of three types of molecular structures:
 - Microtubules
 - Microfilaments
 - Intermediate filaments

Fig. 6-20



Roles of the Cytoskeleton: Support, Motility, and Regulation


- The cytoskeleton helps to support the cell and maintain its shape
- It interacts with **motor proteins** to produce motility
- Inside the cell, vesicles can travel along “monorails” provided by the cytoskeleton
- Recent evidence suggests that the cytoskeleton may help regulate biochemical activities




Components of the Cytoskeleton


- Three main types of fibers make up the cytoskeleton:
 - *Microtubules* are the thickest of the three components of the cytoskeleton
 - *Microfilaments*, also called actin filaments, are the thinnest components
 - *Intermediate filaments* are fibers with diameters in a middle range

Table 6-1



10 μm




10 μm


Actin subunit



 7 nm

5 μm


A horizontal scale bar with vertical end caps, indicating a length of 5 micrometers.

Keratin proteins



**Fibrous subunit (keratins
coiled together)**

8–12 nm

A vertical scale bar with horizontal end caps and arrows at both ends, indicating a thickness of 8–12 nanometers.


Microtubules

- **Microtubules** are hollow rods about 25 nm in diameter and about 200 nm to 25 microns long
- Functions of microtubules:
 - Shaping the cell
 - Guiding movement of organelles
 - Separating chromosomes during cell division

Centrosomes and Centrioles

- In many cells, microtubules grow out from a **centrosome** near the nucleus
- The centrosome is a “microtubule-organizing center”
- In animal cells, the centrosome has a pair of **centrioles**, each with nine triplets of microtubules arranged in a ring

Fig. 6-22



Cilia and Flagella


- Microtubules control the beating of **cilia** and **flagella**, locomotor appendages of some cells
- Cilia and flagella differ in their beating patterns

Video: *Chlamydomonas*

Video: *Paramecium* Cilia

Direction of swimming


(a) Motion of flagella



Direction of organism's movement


Power stroke Recovery stroke

(b) Motion of cilia




-
- Cilia and flagella share a common ultrastructure:
 - A core of microtubules sheathed by the plasma membrane
 - A **basal body** that anchors the cilium or flagellum
 - A motor protein called **dynein**, which drives the bending movements of a cilium or flagellum

Animation: Cilia and Flagella




0.5 μm

(a) Longitudinal section of cilium




(b) Cross section of cilium



(c) Cross section of basal body

-
- How dynein “walking” moves flagella and cilia:
 - Dynein arms alternately grab, move, and release the outer microtubules
 - Protein cross-links limit sliding
 - Forces exerted by dynein arms cause doublets to curve, bending the cilium or flagellum

Fig. 6-25



ATP

— Dynein protein

(a) Effect of unrestrained dynein movement

Cross-linking proteins inside outer doublets


ATP

Anchorage in cell

(b) Effect of cross-linking proteins



(c) Wavelike motion




———— Dynein protein

(a) Effect of unrestrained dynein movement

ATP

**Cross-linking proteins
inside outer doublets**



**Anchorage
in cell**



(b) Effect of cross-linking proteins

1

3


2

(c) Wavelike motion

Microfilaments (Actin Filaments)

- **Microfilaments** are solid rods about 7 nm in diameter, built as a twisted double chain of **actin** subunits
- The structural role of microfilaments is to bear tension, resisting pulling forces within the cell
- They form a 3-D network called the **cortex** just inside the plasma membrane to help support the cell's shape
- Bundles of microfilaments make up the core of microvilli of intestinal cells

Fig. 6-26



-
- Microfilaments that function in cellular motility contain the protein **myosin** in addition to actin
 - In muscle cells, thousands of actin filaments are arranged parallel to one another
 - Thicker filaments composed of myosin interdigitate with the thinner actin fibers

Fig. 6-27

Muscle cell

Actin filament—

Myosin filament—
Myosin arm

(a) Myosin motors in muscle cell contraction

Cortex (outer cytoplasm):
gel with actin network

Inner cytoplasm: sol
with actin subunits

Extending
pseudopodium

(b) Amoeboid movement

Nonmoving cortical
cytoplasm (gel)

Chloroplast

Streaming
cytoplasm
(sol)

Vacuole

Parallel actin
filaments

—Cell wall


(c) Cytoplasmic streaming in plant cells

Muscle cell


Actin filament —

Myosin filament —
Myosin arm /

(a) Myosin motors in muscle cell contraction



(b) Amoeboid movement



(c) Cytoplasmic streaming in plant cells

-
- Localized contraction brought about by actin and myosin also drives amoeboid movement
 - **Pseudopodia** (cellular extensions) extend and contract through the reversible assembly and contraction of actin subunits into microfilaments

-
- **Cytoplasmic streaming** is a circular flow of cytoplasm within cells
 - This streaming speeds distribution of materials within the cell
 - In plant cells, actin-myosin interactions and sol-gel transformations drive cytoplasmic streaming

[Video: Cytoplasmic Streaming](#)

Intermediate Filaments

- **Intermediate filaments** range in diameter from 8–12 nanometers, larger than microfilaments but smaller than microtubules
- They support cell shape and fix organelles in place
- Intermediate filaments are more permanent cytoskeleton fixtures than the other two classes

Concept 6.7: Extracellular components and connections between cells help coordinate cellular activities


- Most cells synthesize and secrete materials that are external to the plasma membrane
- These extracellular structures include:
 - Cell walls of plants
 - The extracellular matrix (ECM) of animal cells
 - Intercellular junctions

Cell Walls of Plants

- The **cell wall** is an extracellular structure that distinguishes plant cells from animal cells
- Prokaryotes, fungi, and some protists also have cell walls
- The cell wall protects the plant cell, maintains its shape, and prevents excessive uptake of water
- Plant cell walls are made of cellulose fibers embedded in other polysaccharides and protein

-
- Plant cell walls may have multiple layers:
 - **Primary cell wall:** relatively thin and flexible
 - **Middle lamella:** thin layer between primary walls of adjacent cells
 - **Secondary cell wall** (in some cells): added between the plasma membrane and the primary cell wall
 - Plasmodesmata are channels between adjacent plant cells

Fig. 6-28



RESULTS

10 μm

**Distribution of cellulose
synthase over time**

**Distribution of microtubules
over time**

The Extracellular Matrix (ECM) of Animal Cells

- Animal cells lack cell walls but are covered by an elaborate **extracellular matrix (ECM)**
- The ECM is made up of glycoproteins such as **collagen, proteoglycans, and fibronectin**
- ECM proteins bind to receptor proteins in the plasma membrane called **integrins**

Fig. 6-30




Fig. 6-30a





Fig. 6-30b



-
- Functions of the ECM:
 - Support
 - Adhesion
 - Movement
 - Regulation


Intercellular Junctions

- Neighboring cells in tissues, organs, or organ systems often adhere, interact, and communicate through direct physical contact
- Intercellular junctions facilitate this contact
- There are several types of intercellular junctions
 - Plasmodesmata
 - Tight junctions
 - Desmosomes
 - Gap junctions

Plasmodesmata in Plant Cells

- **Plasmodesmata** are channels that perforate plant cell walls
- Through plasmodesmata, water and small solutes (and sometimes proteins and RNA) can pass from cell to cell

Fig. 6-31



Tight Junctions, Desmosomes, and Gap Junctions in Animal Cells

- At **tight junctions**, membranes of neighboring cells are pressed together, preventing leakage of extracellular fluid
- **Desmosomes** (anchoring junctions) fasten cells together into strong sheets
- **Gap junctions** (communicating junctions) provide cytoplasmic channels between adjacent cells

Animation: Tight Junctions

Animation: Desmosomes

Animation: Gap Junctions

Fig. 6-32

Tight junctions prevent fluid from moving across a layer of cells





Fig. 6-32a


**Tight junctions prevent
fluid from moving
across a layer of cells**



Tight junction —————

—————
0.5 μm

Desmosome



1 μm

Gap junction —————

—————
0.1 μm

The Cell: A Living Unit Greater Than the Sum of Its Parts

- Cells rely on the integration of structures and organelles in order to function
- For example, a macrophage's ability to destroy bacteria involves the whole cell, coordinating components such as the cytoskeleton, lysosomes, and plasma membrane

Fig. 6-33

5 μm

A vertical scale bar consisting of a central vertical line with short horizontal bars at both ends. To the left of the vertical line, the text "5 μm" is written vertically, reading from bottom to top.

Fig. 6-UN1

	Cell Component	Structure	Function
<p>Concept 6.3 The eukaryotic cell's genetic instructions are housed in the nucleus and carried out by the ribosomes</p>	Nucleus	Surrounded by nuclear envelope (double membrane) perforated by nuclear pores. The nuclear envelope is continuous with the endoplasmic reticulum (ER).	Houses chromosomes, made of chromatin (DNA, the genetic material, and proteins); contains nucleoli, where ribosomal subunits are made. Pores regulate entry and exit of materials.
	Ribosome	Two subunits made of ribosomal RNA and proteins; can be free in cytosol or bound to ER	Protein synthesis
<p>Concept 6.4 The endomembrane system regulates protein traffic and performs metabolic functions in the cell</p>	Endoplasmic reticulum	Extensive network of membrane-bound tubules and sacs; membrane separates lumen from cytosol; continuous with the nuclear envelope.	Smooth ER: synthesis of lipids, metabolism of carbohydrates, Ca ²⁺ storage, detoxification of drugs and poisons Rough ER: Aids in synthesis of secretory and other proteins from bound ribosomes; adds carbohydrates to glycoproteins; produces new membrane
	Golgi apparatus	Stacks of flattened membranous sacs; has polarity (<i>cis</i> and <i>trans</i> faces)	Modification of proteins, carbohydrates on proteins, and phospholipids; synthesis of many polysaccharides; sorting of Golgi products, which are then released in vesicles.
	Lysosome	Membranous sac of hydrolytic enzymes (in animal cells)	Breakdown of ingested substances, cell macromolecules, and damaged organelles for recycling
	Vacuole	Large membrane-bounded vesicle in plants	Digestion, storage, waste disposal, water balance, cell growth, and protection
<p>Concept 6.5 Mitochondria and chloroplasts change energy from one form to another</p>	Mitochondrion	Bounded by double membrane; inner membrane has infoldings (cristae)	Cellular respiration
	Chloroplast	Typically two membranes around fluid stroma, which contains membranous thylakoids stacked into grana (in plants)	Photosynthesis
	Peroxisome	Specialized metabolic compartment bounded by a single membrane	Contains enzymes that transfer hydrogen to water, producing hydrogen peroxide (H ₂ O ₂) as a by-product, which is converted to water by other enzymes in the peroxisome

Fig. 6-UN1a

Concept 6.3


The eukaryotic cell's genetic instructions are housed in the nucleus and carried out by the ribosomes

Cell Component	Structure	Function
Nucleus	Surrounded by nuclear envelope (double membrane) perforated by nuclear pores. The nuclear envelope is continuous with the endoplasmic reticulum (ER).	Houses chromosomes, made of chromatin (DNA, the genetic material, and proteins); contains nucleoli, where ribosomal subunits are made. Pores regulate entry and exit of materials.
Ribosome	Two subunits made of ribosomal RNA and proteins; can be free in cytosol or bound to ER	Protein synthesis

Fig. 6-UN1b

Concept 6.4

The endomembrane system regulates protein traffic and performs metabolic functions in the cell

Cell Component	Structure	Function
Endoplasmic reticulum  (Nuclear envelope)	Extensive network of membrane-bound tubules and sacs; membrane separates lumen from cytosol; continuous with the nuclear envelope.	Smooth ER: synthesis of lipids, metabolism of carbohydrates, Ca ²⁺ storage, detoxification of drugs and poisons Rough ER: Aids in synthesis of secretory and other proteins from bound ribosomes; adds carbohydrates to glycoproteins; produces new membrane
Golgi apparatus	Stacks of flattened membranous sacs; has polarity (<i>cis</i> and <i>trans</i> faces)	Modification of proteins, carbohydrates on proteins, and phospholipids; synthesis of many polysaccharides; sorting of Golgi products, which are then released in vesicles.
Lysosome	Membranous sac of hydrolytic enzymes (in animal cells)	Breakdown of ingested substances cell macromolecules, and damaged organelles for recycling
Vacuole	Large membrane-bounded vesicle in plants	Digestion, storage, waste disposal, water balance, cell growth, and protection

	Cell Component	Structure	Function
Concept 6.5 Mitochondria and chloroplasts change energy from one form to another	Mitochondrion	Bounded by double membrane; inner membrane has infoldings (cristae)	Cellular respiration
	Chloroplast	Typically two membranes around fluid stroma, which contains membranous thylakoids stacked into grana (in plants)	Photosynthesis
	Peroxisome	Specialized metabolic compartment bounded by a single membrane	Contains enzymes that transfer hydrogen to water, producing hydrogen peroxide (H₂O₂) as a by-product, which is converted to water by other enzymes in the peroxisome

You should now be able to:

1. Distinguish between the following pairs of terms: magnification and resolution; prokaryotic and eukaryotic cell; free and bound ribosomes; smooth and rough ER
2. Describe the structure and function of the components of the endomembrane system
3. Briefly explain the role of mitochondria, chloroplasts, and peroxisomes
4. Describe the functions of the cytoskeleton

-
5. Compare the structure and functions of microtubules, microfilaments, and intermediate filaments
 6. Explain how the ultrastructure of cilia and flagella relate to their functions
 7. Describe the structure of a plant cell wall
 8. Describe the structure and roles of the extracellular matrix in animal cells
 9. Describe four different intercellular junctions