

V-HeFT I and V-HeFT II Trials— The Path to A-HeFT

Jay N. Cohn, MD

Professor of Medicine

University of Minnesota Medical School



Major Entry Criteria

V-HeFT I and V-HeFT II

□ Inclusion criteria


- Men, 18 to 75 yr old
- Heart failure ≥ 3 mo
- Reduced exercise capacity (VO_2 max < 25 mL/kg/min)
- Symptomatic despite digitalis and diuretics
- CT ratio > 0.55 , LVEF < 0.45 or LVIDD > 2.7 cm/m²

□ Exclusion criteria

- Hypertension requiring drugs other than diuretics
- Angina requiring frequent or chronic nitrates
- Use of beta-blockers or non-nitrate vasodilators
- Myocardial infarction or cardiac surgery within 3 mo
- Hypertrophic cardiomyopathy or significant valvular disease
- Severe primary lung, liver, or kidney disease

Study Plan

V-HeFT I and V-HeFT II



Study Endpoints


V-HeFT I and V-HeFT II

□ Major endpoints

- All-cause mortality during entire study
- All-cause mortality at 2 yr
- Number and duration of cardiovascular hospitalizations
- Maximum oxygen consumption at peak exercise
- Quality of life (V-HeFT II)

Survival in All Patients

V-HeFT I



Days since randomization date	0	365	730	1095	1460	1825
ISDN/HYD, n =	186	148	109	71	37	16
Placebo, n =	276	202	135	84	41	10
Prazosin, n =	183	135	94	58	27	7

Survival in All Patients

V-HeFT I


V-HeFT I, overall

Treatment	Placebo, n (%)	Drug, n (%)	Hazard ratio (95% CI)	Log-rank P value
ISDN/HYD	120 (44.0)	72 (38.7)	0.78 (0.58, 1.04)	0.093
Prazosin	120 (44.0)	91 (49.7)	1.11 (0.85, 1.46)	0.441

V-HeFT I, endpoint of 2 yr

Placebo, %	ISDN/HYD, %	P value
34.3	25.6	0.053

Survival in All Patients V-HeFT II



ISDN/HYD,	n = 401	332	242	157	86	3
Enalapril,	n = 403	346	265	169	89	1

CV Final NDA20-727_Brief_Document.pdf F.8

Survival in All Patients

V-HeFT II

V-HeFT II, overall

Enalapril n = 403	ISDN/HYD n = 401	Hazard ratio (95% CI)	Log-rank P value
132 (32.8%)	153 (38.2%)	1.23 (0.97, 1.55)	0.083


V-HeFT II, endpoint of 2 yr

Enalapril, %	ISDN/HYD, %	P value
18.0	25.0	0.016

Subgroup Analysis

Survival in Black Patients and White Patients

V-HeFT II





Patients, n

ISDN/HYD	109	92	67	49	29	1	282	231	171	105	55	1
Enalapril	106	93	69	47	24	2	292	251	194	123	66	1

Survival in Black Patients and White Patients

V-HeFT I



Patients, n

ISDN/HYD	49	43	36	28	16	8	132	102	71	42	22	9
Placebo	79	61	44	29	14	3	192	140	91	55	27	8

V-HeFT I—Conclusions (1)

- ISDN/HYD compared to placebo was associated with
 - A 22% lower risk of death overall ($P = 0.09$)
 - A 12% lower risk of death in white patients ($P = 0.47$)
 - A 47% lower risk of death in black patients ($P = 0.04$)

V-HeFT II—Conclusions (2)

- Enalapril compared to ISDN/HYD was associated with
 - A 23% lower mortality overall ($P = 0.08$)
 - A 39% lower mortality in white patients ($P = 0.02$)
 - No difference in mortality in blacks

From V-HeFT I and V-HeFT II to A-HeFT

- Based on V-HeFT I and V-HeFT II, a clinical study was needed to confirm the hypothesis that the ISDN/HYD combination benefits outcomes in black HF patients
- A-HeFT was designed as a prospective, placebo-controlled study with the objective of testing BiDil's effects on survival, heart failure hospitalizations, and quality of life in patients receiving contemporary therapy for heart failure