

volumes


Polyhedrons



What is a polyhedron?




These are polyhedrons.



These are not polyhedrons.


Circles are not polygons


Identifying Polyhedrons



- A polyhedron is a solid that is bounded by polygons, called *faces*, that enclose a single region of space.
- An edge of polyhedron is a line segment formed by the intersection of two faces
- A vertex of a polyhedron is a point where three or more edges meet

Parts of a Polyhedron








Example 1

Counting Faces, Vertices, and Edges

 Count the faces, vertices, and edges of each polyhedron








Example 1A Counting Faces


Count the faces, vertices, and edges of each polyhedron



4 faces

Example 1a Counting Vertices


Count the faces, vertices, and edges of each polyhedron



4 vertices

Example 1a Counting Edges


Count the faces, vertices, and edges of each polyhedron



6 edges

Example 1b Counting Faces


Count the faces, vertices, and edges of each polyhedron



5 faces

Example 1b Counting Vertices


Count the faces, vertices, and edges of each polyhedron



5 vertices

Example 1b Counting Vertices


Count the faces, vertices, and edges of each polyhedron



8 edges


Example 1c Counting Faces

Count the faces, vertices, and edges of each polyhedron



Example 1c Counting Vertices


Count the faces, vertices, and edges of each polyhedron



6 vertices

Example 1c Counting Edges

Count the faces, vertices, and edges of each polyhedron



10 edges


Notice a Pattern?

Faces	Vertices	Edges
4	4	6
5	5	8
6	6	10


Theorem 12.1 Euler's Theorem

The number of faces (F), vertices (V), and edges (E) of a polyhedron is related by F + V = E + 2

- More about Polyhedrons
 The surface of a polyhedron consists of all points on its faces
 - A polyhedron is convex if any two points on its surface can be connected by a line segment that lies entirely inside or on the polyhedron




These polyhedrons are convex.




These polyhedrons are not convex.

Regular Polyhedrons

 A polyhedron is regular if all its faces are congruent regular polygons.




5 kinds of Regular Polyhedrons




Regular tetrahedron

4 faces




Cube

6 faces



Regular octahedron

8 faces



Regular dodecahedron


Regular icosahedron

12 faces

20 faces


Example 2 Classifying Polyhedrons

One of the octahedrons is regular. Which is it?




A polyhedron is **regular** if all its faces are congruent regular polygons.


Example 2 Classifying Polyhedrons



All its faces are congruent equilateral triangles, and each vertex is formed by the intersection of 4 faces



Faces are not all congruent (regular hexagons and squares)



Faces are not all regular polygons or congruent (trapezoids and triangles)

Example 3

Counting the Vertices of a Soccer Ball

A soccer ball has 32 faces: 20 are regular hexagons and 12 are regular pentagons. How many vertices does it have?




 A soccer ball is an example of a semiregular polyhedron - one whose faces are more than one type of regular polygon and whose vertices are all exactly the same

Example 3 Counting the Vertices of a Soccer Ball

- A soccer ball has 32 faces: 20 are regular hexagons and 12 are regular pentagons. How many vertices does it have?
- Hexagon = 6 sides, Pentagon = 5 sides
- Each edge of the soccer ball is shared by two sides
- Total number of edges = $\frac{1}{2}(6 \cdot 20 + 5 \cdot 12) = \frac{1}{2}(180) = 90$
- Now use Euler's Theorem
- F + V = E + 2
- 32 + V = 90 + 2
- V = 60


Prisms

- A prism is a polyhedron that has two parallel, congruent faces called bases.
- The other faces, called lateral faces, are parallelograms and are formed by connecting corresponding vertices of the bases
- The segment connecting these corresponding vertices are lateral edges
- Prisms are classified by their bases




Prisms

- The altitude or height, of a prism is the perpendicular distance between its bases
- In a right prism, each lateral edge is perpendicular to both bases
- Prisms that have lateral edges that are oblique (#90°) to the bases are oblique prisms
- The length of the oblique lateral edges is the slant height of the prism



Right rectangular prism



Oblique triangular prism

Surface Area of a Prism

 The surface area of a polyhedron is the sum of the areas of its faces


Example 1 Find the Surface Area of a Prism

The Skyscraper is 414 meters high. The base is a square with sides that are 64 meters. What is the surface area of the skyscraper?



Example 1 Find the Surface Area of a Prism

The Skyscraper is 414 meters high. The base is a square with sides that are 64 meters. What is the surface area of the skyscraper?



Surface Area = $4(64 \cdot 414) + 2(64 \cdot 64) = 114,176 \text{ m}^2$

Example 1 Find the Surface Area of a Prism


The Skyscraper is 414 meters high. The base is a square with sides that are 64 meters. What is the surface area of the skyscraper?

Surface Area = $4(64 \cdot 414) + 2(64 \cdot 64) = 114,176 \text{ m}^2$

Surface Area = $(4 \cdot 64) \cdot 414 + 2(64 \cdot 64) = 114,176 \text{ m}^2$


height


Perimeter of Area of the base



Nets

A net is a pattern that can be cut and folded to form a polyhedron.





Surface Area of a Right Prism


The surface area, S, of a right prism is


$$S = 2B + Ph$$

where B is the area of a base, P is the perimeter of a base, and h is the height

Example 2 Finding the Surface Area of a Prism

Find the surface area of each right prism






Example 2

Finding the Surface Area of a Prism

Find the surface area of each right prism



$$S = 2B + Ph$$

Area of the Base = 5x12=60

Perimeter of Base = 5+12+5+12 = 34

Height of Prism = 8


$$\bullet S = 2B + Ph$$

$$\bullet$$
S = 2(60) +(34) \bullet 8

$$\bullet$$
S = 120 + 272 = 392 in²

Example 2 Finding the Surface Area of a Prism

Find the surface area of each right prism



$$S = 2B + Ph$$

Area of the Base = $\frac{1}{2}(5)(12)=30$

Perimeter of Base = 5+12+13=30

Height of Prism = 4 (distance between triangles)


$$\bullet S = 2B + Ph$$

$$\bullet$$
S = 2(30) +(30) \bullet 4

$$\bullet$$
S = 60 + 120 = 180 in²

Cylinders

- A cylinder is a solid with congruent circular bases that lie in parallel planes
- The altitude, or height, of a cylinder is the perpendicular distance between its bases
- The lateral area of a cylinder is the area of its curved lateral surface.
- A cylinder is right if the segment joining the centers of its bases is perpendicular to its bases



The surface area of a right cylinder is the sum of the areas of its two bases and its lateral area.

Surface Area of a Right Cylinder


The surface area, S, of a right circular cylinder is

$$S = 2B + Ch$$
or
$$2\pi r^2 + 2\pi rh$$

where B is the area of a base, C is the circumterence of a base, r is the radius of a base, and h is the height

Example 3 Finding the Surface Area of a Cylinder

Find the surface area of the cylinder



Example 3


Finding the Surface Area of a Cylinder

- Find the surface area of the cylinder
- \Box $2\pi r^2 + 2\pi rh$
- \Box 2 π (3)² + 2 π (3)(4)
- \Box 18 π + 24 π
- $\Box 42\pi \approx 131.9 \text{ ft}^2$

L


Radius = 3

Height = 4




Pyramids


A pyramid is a polyhedron in which the base is a polygon and the lateral faces are triangles that have a common vertex



Pyramids

- The intersection of two lateral faces is a lateral edge
- The intersection of the base and a lateral face is a base edge
- The altitude or height of the pyramid is the perpendicular distance between the base and the vertex






Regular Pyramid

 A pyramid is regular if its base is a regular polygon and if the segment from the vertex to the center of the base is perpendicular to the base

 The slant height of a regular pyramid is the altitude of any lateral face (a nonregular


pyramid has no slant height)



Regular pyramid

Developing the formula for surface area of a regular pyramid

- Area of each triangle is ½bL
- Perimeter of the base is 6b
- Surface Area =(Area of base) + 6(Area of lateral faces)
- $S = B + 6(\frac{1}{2}b)$
- $S = B + \frac{1}{2}(6b)(1)$
- $S = B + \frac{1}{2}P$



Surface Area of a Regular Pyramid


The surface area, S, of a regular pyramid is


$$S = B + \frac{1}{2}P$$

Where B is the area of the base, P is the perimeter of the base, and L is the slant height

Example 1 Finding the Surface Area of a Pyramid

 Find the surface area of each regular pyramid






Example 1 Finding the Surface Area of a Pyramid

Find the surface area of each regular

a.



$$S = 25 + \frac{1}{2}(20)(4)$$

= 25 + 40
= 65 ft²

Base is a Square

Area of Base = 5(5) = 25

Perimeter of Base


$$5+5+5+5=20$$

Slant Height = 4

Example 1 Finding the Surface Area of a Pyramid

Find the surface area of each regular pyramid

$$S = B + \frac{1}{2}PL$$



$$S = 54\sqrt{3} + \frac{1}{2}(36)(8)$$
$$= 54\sqrt{3} + 144$$
$$= 237.5 \text{ m}^2$$

Base is a Hexagon A=1/2aP


$$A \square \frac{1}{2} (3\sqrt{3})(36) \square 54\sqrt{3}$$

Perimeter = 6(6)=36

Slant Height = 8

Cones

- A cone is a solid that has a circular base and a vertex that is not in the same plane as the base
- The lateral surface consists of all segments that connect the vertex with point on the edge of the base
- The altitude, or height, of a cone is the perpendicular distance between the vertex and the plane that contains the base



Right Cone

- A right cone is one in which the vertex lies directly above the center of the base
- The slant height of a right cone is the distance between the vertex and a point on the edge of

Slant height

Base

Height


Right cone

Lateral

surface

base

Developing the formula for the surface area of a right cone



- Use the formula for surface area of a pyramid S = B + ½P/
- As the number of sides on the base increase it becomes nearly circular
- Replace ½P (half the perimeter of the pyramids base) with πr (half the circumference of the cone's base)

Surface Area of a Right Cone


The surface area, S, of a right cone is

$$S = \pi r^2 + \pi r \ell$$

Where r is the radius of the base and L is the slant height of the cone


Example 2 Finding the Surface Area of a Right Cone

Find the surface area of the right cone



Example 2 Finding the Surface Area of a Right Cone

Find the surface area of the right cone



$$S = \pi r^{2} + \pi r/$$

$$= \pi (5)^{2} + \pi (5)(7)$$

$$= 25\pi + 35\pi$$

$$= 60\pi \text{ or } 188.5 \text{ in}^{2}$$

Radius = 5
Slant height = 7

Volume formulas

The Volume, V, of a prism is V = BhThe Volume, V, of a cylinder is $V = \pi r^2 h$

The Volume, V, of a pyramid is $V = \frac{1}{3}Bh$ The Volume, V, of a cone is $V = \frac{1}{3}\pi r^2h$

The Surface Area, S, of a sphere is $S = 4\pi r^2$ The Volume, V, of a sphere is $V = \frac{4}{3}\pi r^3$

Volume

 The volume of a polyhedron is the number of cubic units contained in its interior

· Label volumes in cubic units like cm³.

in³, ft³, etc


Postulates

- All the formulas for the volumes of polyhedrons are based on the following three postulates
 - Volume of Cube Postulate: The volume of a cube is the cube of the length of its side, or $V = s^3$
 - Volume Congruence Postulate: If two polyhedrons are congruent, then they have the same volume
 - Volume Addition Postulate: The volume of a solid is the sum of the volumes of all its nonoverlapping parts

Example 1: Finding the Volume of a Rectangular Prism


The cardboard box is 5" x 3" x 4"
How many unit cubes can be packed
into the box? What is the volume of

the box?



Example 1: Finding the Volume of a rectangular Prism

• The cardboard box is $5" \times 3" \times 4"$ How many unit cubes can be packed into the box? What is the volume of the box?




- ·How many cubes in bottom layer?
- $\cdot 5(3) = 15$
- ·How many layers?
- •4
- \cdot V=5(3)(4) = 60 in³

 $V = L \times W \times H$ for a rectangular prism


Volume of a Prism and a Cylinder

Cavalieri's Principle

If two solids have the same height and the same cross-sectional area at every level, then they have the same volume



Because each stack has the same number of pennies, it follows that each stack has the same volume.



Volume of a Prism

· The Volume, V, of a prism is

where B is the area of a base and h is the height

Volume of a Cylinder


The volume, V, of a cylinder is V=Bh or

$$V = \pi r^2 h$$

where B is the area of a base, h is the height and r is the radius of a base

Example 2 Finding Volumes


 Find the volume of the right prism and the right cylinder



Example 2 Finding Volumes

 Find the volume of the right prism and the right cylinder

a.



Area of Base

$$B = \frac{1}{2}(3)(4) = 6$$

Height = 2

$$V = Bh$$

3 cm

$$V = 6(2)$$


4 cm

$$V = 12 \text{ cm}^3$$

Example 2 Finding Volumes

 Find the volume of the right prism and the right cylinder

b.



Area of Base

$$B = \pi(7)^2 = 49\pi$$


$$Height = 5$$

$$V = 49\pi(5)$$

$$V = 245\pi \text{ in}^3$$

Example 3 Estimating the Cost of Moving

 You are moving from Newark, New Jersey, to Golden, Colorado - a trip of 2000 miles. Your furniture and other belongings will fill half the truck trailer. The moving company estimates that your belongings weigh an average of 6.5 pounds per cubic foot. The company charges \$600 to ship 1000 pounds. Estimate the cost of shipping your belongings.



Example 3 Estimating the Cost of Moving

 You are moving from Newark, New Jersey, to Golden, Colorado - a trip of 2000 miles. Your furniture and other belongings will fill half the truck trailer. The moving company estimates that your belongings weigh an average of 6.5 pounds per cubic foot. The company charges \$600 to ship 1000 pounds. Estimate the cost of shipping your belongings.

Volume = $L \times W \times H$

V = 50(8)(9)

 $V = 3600 \text{ ft}^3$

 $3600 \div 2 = 1800 \text{ ft}^3$

1800(6.5) = 11,700 pounds

 $11,700 \div 1000 = 11.7$

11.7(600) = \$7020

