

Attention-deficit/hyperactivity disorder (ADHD) is a brain disorder marked by an ongoing pattern of inattention and/or hyperactivity-impulsivity that interferes with functioning or development.



Inattention means a person

- wanders off task,
- lacks persistence,
- has difficulty sustaining focus,
- is disorganized;


These problems are not due to lack of comprehension.



Hyperactivity means a person


- seems to move about constantly, including in situations in which it is not appropriate;
- excessively fidgets, taps, or talks.

In adults, it may be extreme restlessness.




Impulsivity means a person

- makes hasty actions that occur in the moment without first thinking about them;
- •a desire for immediate rewards or inability to delay gratification.




Attention deficit hyperactivity disorder (ADHD)


- •is a dysfunction of the Reticular Activating System (RAS).
- •RAS is the center of consciousness that
- 1.coordinates learning and memory,
- 2.and which normally supplies the appropriate neural connections necessary for smooth information processing and clear, non-stressful attention.



- The RAS appears to be intimately involved in
- the neural mechanisms which produce consciousness and focused attention
- •It receives impulses from the spinal cord and relays them to the Thalamus, and from there to the Cortex, and back again in a feedback loop to the Hippocampus/Thalamus/Hypothalamus



- Without continual excitation of cortical neurons by reticular activation impulses, an individual is unconscious and cannot be aroused.
- When stimulation is enough for consciousness but not for attentiveness, ADD results.
- If too activated, an individual cannot relax or concentrate (and is over-stimulated or hyperactive) often resulting in ADHD.



Risk Factors

Scientists are not sure what causes ADHD.

A number of factors can contribute to ADHD, such as:

- Genes
- Cigarette smoking, alcohol use, or drug use during pregnancy
- Exposure to environmental toxins during pregnancy
- •Exposure to environmental toxins, such as high levels of lead, at a young age
- Low birth weight
- Brain injuries

- ADHD is more common in males than females,
- Females with ADHD are more likely to have problems primarily with inattention.
- Other conditions, such as learning disabilities, anxiety disorder, conduct disorder, depression, and substance abuse, are common in people with ADHD.

