

# Phylum Mollusca



























Possess trochophore larvae

Mollusca and Annelida are closely allied phyla

93,000 species of described molluscs (extant) +  
70,000 more species from fossil record





# Defining characteristics of Mollusca

- Bilateral symmetry, cephalization
- Coelom (around heart)
- Mantle
  - draped over viscera
  - secretes shell
- Complete digestive tract, Radula
- Trochophore larvae (often also veliger larvae)
- Locomotion by muscular foot
- Heart, liver, gills (ctenidia), kidney

*Phylum.* Mollusca

*Class.* Aplacophora

*Class.* Polyplacophora

*Class.* Monoplacophora


*Class.* Gastropoda

*Class.* Cephalopoda


*Class.* Bivalvia

*Class.* Scaphopoda









# Mollusc origins?




**Platyhelminthe-like ancestor with spicules on dorsal surface, cilia on ventral surface, and dorsoventral musculature.**







# Mollusc origins?




**Spicules fuse to form dorsal shell, cilia on ventral surface, and dorsoventral musculature.**








# Mollusc origins?



**Spicules fuse to form dorsal shell, cilia on ventral surface, dorsoventral musculature, gonads and excretory pores in mantle cavity**




# Mollusc origins?



**Spicules fuse to form dorsal shell, cilia on ventral surface, dorsoventral musculature, gonads, excretory pores, and ctenidia, in mantle cavity.**


# HAM: Hypothetical ancestral mollusc






Dorsal mantle covers the visceral mass.

Secretes the shell




# Mollusc shell


- Periostracum
- Prismatic layer
- Nacreous layer
- Mantle epithelium
- Mantle lobes



# Pearl formation




# Ctenidium (Respiration)







# Ctenidium (gill)




# Paired ventral nerve cords



# Complete digestive system





# Radula





# Radula




BIODIDAC © Houseman, Univ. of Ottawa



BIODIDAC © Houseman, Univ. of Ottawa

# Aplacophora



# Class Aplacophora



- 320 spp, all marine
- No fossil record (!) & poorly studied
- Calcareous spines and scales in epidermis (no true shell)
- Radula used for grasping
- Burrow in the substrate
- Eat cnidarians







# Polyplacophora




# Polyplacophora





# Polyplacophora



# Polyplacophora

- The “chitons”
- 800 spp, marine
- Typically, shell = 8 dorsal plates
- Ctenidia use counter-current gas exchange
- Commonly encountered in the intertidal zone

# Class Gastropoda



- Single external shell
- Radula for scraping food
- Moves in wave like contractions through slime
- Can be hermaphrodites, though commonly not
- Ex) Snails, slugs



# Class Gastropoda

- Snails live on land or in fresh or salt water
  - They have eyes on tentacles on their head
- Slugs live on land and Sea Slugs (a.k.a. nudibranches) live in the ocean
  - Slugs do not have shells
  - Have exchange of oxygen (diffusion) across their entire body

# Gastropoda

- More active than mono and polyplacophorans
  - Highly cephalized: tentacles, eyes
- Gonochoristic
- Veliger larva (an advanced version of the trochophore larva)



# Class Gastropoda

Three evolutionary innovations occurred among the gastropods: changes in the shell, increased development of the head, the embryonic process of torsion


## 1. Changes in the Shell

The shell became higher and conical with a reduced aperture

The shell also became coiled

Shells initially were **planospiral** - bilaterally symmetrical shell with the whorls lying in the same plane



Modern day shells are **asymmetrical** - each successive coil is a little outside and offset a little above the one below



# Shell

Most have a single, spiraled shell and can move the entire head and foot into this shell for protection.




Also, many gastropods have a hardened plate called the **operculum** on the back of the foot that plugs the shell aperture when the body is withdrawn



# Nutrition

Many gastropods are herbivores and use their radula scrap algae from surfaces of rocks

Some gastropods are active predators and in these the radula is often highly modified, e.g., as a drill (oyster drills) or harpoon (venomous gastropods)




Cone snail

# Respiration

Aquatic gastropods possess gills for respiration

Terrestrial gastropods obtain oxygen via a well vascularized mantle



gills




Vascularized  
mantle



# Pulmonata

- Highly vascularized mantle for gas exchange (lung)
- 17,000 spp: slugs, pond snails





# Sea Slugs!



# Sea Slugs!





# Mollusks (Phylum Mollusca)

- **Bivalvia – clam:**
  - **two valves, connected by dorsal hinge ligament**
  - **adductor muscles (used by living clam to close the shell)**
  - **gills in mantle cavity**
  - **wedgelike foot**





# Class Bivalvia – Two Shells

- Three layers make up the shell of a bivalve
  - Inner most protects the body of the animal
  - Middle layer strengthens the shell with calcium carbonate
  - Outer layer protects against acid in the water

# Class Bivalvia


Shells divided into 2 equal halves or **valves**  
Mantle tissue is indented in the anterior-posterior margins, with 2 centers of calcification

Shells joined at the dorsal midline by a non calcified protein ligaments called the **hinge**






# Movement of the Ventilating Currents



# Adaptive Radiation of Bivalves

Chapter



**Figure 29.35**

Bivalve life styles. All of these species would not found together in the same



# Reproduction


Most are dioecious

Marine forms usually produce free swimming trochophore and veliger larvae

Many of the freshwater bivalves have a different life history pattern; produce larvae called **glochidia**

Glochidia are housed in the outer gills; they use there outer gill as a brood chamber - **marsupium**

When the glochidia are released they parasitize the fins and gills of fishes



# **Mollusks (Phylum Mollusca)**

**:**

- **Cephalopoda – squid:**

  - muscular foot with tentacles (with suction discs)**

  - thick mantle but no external shell (true for most cephalopods)**

- **excurrent siphon (for jet propulsion)**

  - large eyes**

# Class Cephalopoda




- Means “head foot”
- Includes octopuses, squids, cuttlefishes and chambered nautiluses
- Free swimming and predatory
- Tentacles with suction cups to grasp prey
- Have jaws called beaks to destroy their prey
- Has the largest invertebrate brain



# Cephalopods

- Can learn to perform tasks
- Has complex eyes
- Closed circulatory system
- Are sexual - internal
- Have ink to confuse predators



# Cephalopoda

- Swift, agile carnivores
- Closed circulatory system, 2 hearts
- Separate sexes
- Foot modified to form arms, tentacles, siphon
- Brain, cranium, complex image-forming eye
- 700 extant spp, 10,000 extinct spp
- Arose from limpet-like monoplacophorans
- Ergo, ventral became function anterior, etc

# Cephalopods



**School Bus**  
40 ft (12 m)



**Giant Squid**  
33 ft (10 m)  
Avg. Est. Length



**Colossal Squid**  
43 ft (13 m)  
Max. Est. Length  
Avg. Length Unknown

# Cephalopoda

- Endocochleate cephalopods
  - Reduce internal shell, or shell absent
  - Squids, cuttlefish, octopi




# Locomotion

- Cephalopods are excellent swimmers: streamlined body; tentacles and fins as stabilizers
- Swim by means of jet propulsion, using the highly modified muscular mantle and the **siphon**

By relaxing the mantle the mantle cavity is expanded and water can be drawn in

By contracting the mantle water can be forced out of the mantle cavity by means of the small siphonal opening







# Other General Features

For protection, they possess an **ink sacs**

Cephalopods have well-developed sense organs, including a camera type eye

Some have well-developed brains and show a remarkable capacity for learning.

Cephalopods are the only molluscan class with a **closed circulatory system**




FIGURE 9-33

Eye of a cuttlefish (*Sepia*). The structure of cephalopod eyes shows a high degree of convergent evolution with the eyes of vertebrates.






Photo: Dave Knight











# Cephalopoda


Posterior surface


Right

Ventral


Dorsal

Left





**Shell (Pen)**



**Systemic heart**


**Branchial heart**

**Ctenidium**

**Funnel**

Hectocotylus (sperm-bearing arm in males)

**Reproduction:** trochophore and veliger are bypassed and hatch into planktonic juveniles





Brain is surrounded by a cranium

9/9/97



# Cephalopod eye










# Examples of Cephalopods



A giant squid (3.15-metre-long) has netted off the UK coast; first time in 15 years.

The squid, believed to be female and three years old, did not survive being brought to the surface.





# The Mimic Octopus



PHOTOS BY ROGER STEENE

**EVEN A SEA SNAKE:** Not only can a mimic octopus change its shape to imitate a sea snake (a real version is at right), it can also quickly change its coloration to match by using 'color sacs' and 'reflectors' in its skin.



# Cephalopoda

- Ectocochleate cephalopods
  - Have external shell with internally subdivisions used for buoyancy control
  - This ancestral group is almost completely extinct
  - E.g. *Nautilus*



Nautilus is the only cephalopod with an external shell and lacking chromatophores

Chromatophores (color cells)

Iridocytes (reflective cells)

- Millions of these allow rapid changes in color, polarized signals
- Also have photophores for bioluminescence

Cephalopods except Nautilus have ink sac