


Amines



Amines


Properties of amines

Synthesis of amines


Reactions of amines

Summary activities


Amines are nitrogen-containing organic compounds derived from ammonia, where one or more of the hydrogen atoms has been replaced by an **alkyl** or **aryl** group.



ammonia



methylamine




phenylamine


Amines have unpleasant odours: those with low boiling points smell like ammonia, whereas those that are liquid at room temperature have fishy aromas.

An alkyl or aryl group can be represented by an R when drawing a chemical structure. This is referred to as an **R group**.


- **Primary (1°) amines** have one R group attached to the nitrogen atom.



- **Secondary (2°) amines** have two R groups attached to the nitrogen atom.





- **Tertiary (3°) amines** have three R groups attached to the nitrogen atom.




Shape of amines

The ammonium ion (NH_4^+) is **tetrahedral** in shape, as the four bonding pairs of electrons (which repel each other) spread out equally around the central nitrogen atom.



Ammonia (NH_3) is **pyramidal** in shape, as it has a lone pair in place of one bonding pair, which exerts a stronger repulsive force.

The amines are shaped similarly to ammonia, with a bond angle of 107° between groups on the nitrogen atom.



Do these structures
represent primary,
secondary or tertiary
amines?


Press **start** to begin.

start




Naming amines (1/3)


Amines are named using the suffix *-amine*.



methylamine




ethylamine




propylamine

If two identical R groups are attached, the prefix *di-* is used, and if three identical groups are present, then *tri-* is used.




diethylamine




triethylamine

Naming amines (2/3)


If two different alkyl or aryl groups are present, they are listed alphabetically.




ethyl**methyl**amine



ethyl**dipropyl**amine




methyl**phenyl**amine




methyl**diphenyl**amine

Naming amines (3/3)


If other functional groups are present in the molecule, the presence of amine groups is denoted using the *amino-* prefix.



2-aminoethanoic acid




3-aminopropanoic acid



1,2-diaminopentane

What's the name of the amine?

Structure: 1/6




*type in the name of
the structure below*



Solubility of primary amines

Shorter chain amines are soluble in water, as the lone pair of electrons on the nitrogen atom allows them to form hydrogen bonds with water molecules.

Longer chain amines are only sparingly soluble, as the larger R groups interfere with the hydrogen bonds.



In aqueous solution, amine molecules are able to accept an H^+ ion from the water molecules, resulting in an alkaline solution due to the remaining OH^- ions.

Boiling points of primary amines

It is useful to compare the boiling point of methylamine (CH_3NH_2) with that of ethane (CH_3CH_3) because both molecules contain the same number of electrons and have roughly the same shape.


Boiling point ($^{\circ}\text{C}$)			
alkane		primary amine	
CH_3CH_3	-88.6	CH_3NH_2	-6.3
$\text{CH}_3\text{CH}_2\text{CH}_3$	-42.0	$\text{CH}_3\text{CH}_2\text{NH}_2$	16.6
$\text{CH}_3(\text{CH}_2)_2\text{CH}_3$	-0.5	$\text{CH}_3(\text{CH}_2)_2\text{NH}_2$	48.6

The main reason the boiling points of the primary amines are higher is that they can form hydrogen bonds with each other.



Boiling points of isomeric amines

Secondary amines also form hydrogen bonds, but as the nitrogen atom is in the middle of the hydrocarbon chain the strength of the dipole is slightly less. This decreases the strength of dipole–dipole attractions.



amine type	formula	boiling point (°C)
1°	$\text{CH}_3(\text{CH}_2)_2\text{NH}_2$	48
2°	$\text{CH}_3\text{CH}_2\text{NHCH}_3$	37
3°	$(\text{CH}_3)_3\text{N}$	3

In tertiary amines, there are no hydrogens attached directly to the nitrogen, so hydrogen bonding between molecules is impossible.



Amines

Properties of amines


Synthesis of amines

Reactions of amines

Summary activities




Halogenoalkanes will undergo **nucleophilic substitution** reactions with **ethanolic ammonia** to form a primary amine:



The primary amine may then nucleophilically attack another molecule of halogenoalkane, to form a secondary amine:




A tertiary amine can be formed by nucleophilic attack of a halogenoalkane by a secondary amine:





In reality, a mixture of the above products is usually formed, which must be separated by distillation.

Nitriles can be reduced to primary amines using hydrogen in the presence of a nickel catalyst:




Nitriles can also be reduced to primary amines using strong reducing agents such as lithium tetrahydridoaluminate (LiAlH_4), which can be represented as $[\text{H}]$:



Preparation of phenylamine


Aromatic amines can be prepared by the reduction of nitrated arenes using a mixture of tin metal and concentrated hydrochloric acid:




This method is commonly used to prepare aromatic amines in the lab. A similar method, using iron instead of tin, is used to prepare phenylamine industrially.



Which conditions?



A large white rectangular area with a dark blue border, intended for writing or drawing.



A toolbar at the bottom of the white area containing icons for an eraser, a yellow highlighter, a black pen, a left arrow, a right arrow, an undo button, and a redo button.



Amines


Properties of amines

Synthesis of amines

Reactions of amines


Summary activities

Aliphatic amines have at least one alkyl group bonded to the nitrogen. The lone pair of electrons on the nitrogen means that aliphatic amines behave similarly to ammonia:



- they act as **nucleophiles** and take part in reactions involving donation of the lone pair
- they act as **Brønsted–Lowry bases** (H^+ acceptors).

Aromatic amines contain a benzene ring directly attached to the NH_2 group.




The delocalized system of the benzene group is able to incorporate the lone pair of electrons from the nitrogen atom, meaning that aromatic amines have different properties to aliphatic amines.

Amines as Brønsted–Lowry bases

The lone pair of electrons on the nitrogen atom means that amines can act as **Brønsted–Lowry bases**.

Press **play** to find out more.



Relative base strength

Aliphatic amines are stronger bases (lower pK_b) than ammonia.


This is because alkyl groups repel electrons, leading to an increase in negative charge around the nitrogen so that it more readily attracts and accepts an H^+ ion. This means, 2° amines are more basic than 1° amines, and 3° amines are more basic still.

Compound	pK_b
NH_3	4.75
CH_3NH_2	3.36
$(CH_3)_2NH$	3.27
$C_6H_5NH_2$	9.38

Phenylamine is less basic than ammonia because the phenyl ring is an **electron-withdrawing** group. The lone pair of electrons interact with the delocalized electrons in the ring, and so are less readily donated to an H^+ ion.

Reactions of amines as bases


Amines accept protons (H^+) from acids to form salts:



ethylamine

ethylammonium chloride


If the reaction is carried out in solution, the amine accepts an H^+ from a hydroxonium ion to form an ionic salt and water (a neutralization reaction):



Amines as nucleophiles

Aliphatic amines act as **nucleophiles**, attacking and substituting the halogen atom in a halogenoalkane.


Press **play** to find out how this occurs.



Uses of quaternary ammonium salts

Quaternary ammonium salts are salts of a quaternary ammonium cation (NR_4^+) and an anion.

They are used as **cationic surfactants** in products such as fabric conditioner and shampoo. Their purpose is to smooth fabric or hair, making them softer, by reducing surface tension.



Other quaternary ammonium salts, similar to those shown *left*, are used as **antiseptics** in products such as wet wipes.

 $n = 8, 10, 12, 14, 16, 18$



Reaction of amines with acyl compounds

Amines are able to act as nucleophiles, and react with **acyl compounds** (acid chlorides and acid anhydrides) via an addition–elimination mechanism.

Press the buttons to find out more about how this occurs


reaction with
acid chlorides

reaction with
acid anhydrides



Synthesis of diazonium salts

An aromatic amine can be reacted with nitrous acid (HNO_2) to produce a diazonium ion:




**benzenediazonium
chloride**

The diazonium ion is very unstable, so the temperature of the reaction mixture must be kept below 10°C in order to prevent the ion from decomposing.

Coupling reactions

The positive charge of the nitrogen makes the diazonium ion an electrophile, which is able to undergo electrophilic substitution with a benzene ring.




This is an example of a **coupling reaction**. It must take place under alkaline conditions.

The product is an **azo compound**.

Azo compounds are highly coloured. They are also stable and resistant to fading, and so are important in the **dye** industry.

The colour results from the joining of the two delocalized electron systems via the N=N group. Colour can be altered by changing the number and type of functional groups attached.



methyl orange

Methyl orange indicator is an azo compound. The colour of the molecule changes when H^+ ions are added across the N=N bond.

Reactions of amines: true or false?

Are these statements about reactions of amines true or false?

1.	Phenylamines are more basic than alkyl amines.	<input data-bbox="1508 315 1808 394" type="text" value="?"/>
2.	Some quarternary ammonium salts can be used as antiseptics.	<input data-bbox="1508 468 1808 546" type="text" value="?"/>
3.	Amines can react with acyl chlorides to form a substituted amide.	<input data-bbox="1508 621 1808 699" type="text" value="?"/>
4.	Aromatic amines are refluxed with nitrous acid to form azo compounds.	<input data-bbox="1508 773 1808 852" type="text" value="?"/>
5.	Phenol is used in coupling reactions as the OH group is electron-withdrawing from the benzene ring.	<input data-bbox="1508 926 1808 1005" type="text" value="?"/>

true

false



Amines

Properties of amines

Synthesis of amines

Reactions of amines

Summary activities

Glossary of keywords: amines

acid anhydride – Symmetrical acid prepared by the reaction of two molecules of carboxylic acid, resulting in the elimination of a water molecule.

acid chloride – A compound that contains an **acyl group** with a chlorine attached to the acyl carbon ($R(CO)Cl$). Also known as an acyl chloride.

acyl group – The $RC=O$ group found in esters, **acid chlorides** and **acid anhydrides**.

addition-elimination – Mechanism by which

ABC

DEF

GHI

JKL

MNO

PQR

STU

VWX

YZ



How good is your knowledge
of keywords relating
to **amines**?

Press **start** to find out.

start



How much do you know
about **amines** and
their reactions?

Press **start** to begin this
multiple choice quiz.

start

