

Toshkent Davlat Stomatologiya Instituti
Bolalar Stomatologiyasi fakulteti

307 “c” guruh talabasi

Pulatov Avazning

Dermatologiya fanidan

MUSTAQIL

ISHI

Mavzu: Yiringli kasalliklar

Yuza	Chuqur
I. Stafilodermiyalar	
<ol style="list-style-type: none"> 1. Osteofollikulit 2. Uza follikulit 3. Oddiy sikoz 4. Oddiy xusnbuzar 5. Gudaklarni epidemik porsildoq yarasi 	<ol style="list-style-type: none"> 1. Chuqur follikulit 2. Furunkul, furunkulez 3. karbunkul 4. gidradenit
II. streptodermiyalar	
<p>. Streptokokkli impetigo</p> <ol style="list-style-type: none"> A) Streptokokkli bichilish B) intertriginoz B) bullez Г) xalqasimon Д) zahmga uxshash Г) yuza panariciy <p>2. Quruq streptodermiya</p> <ol style="list-style-type: none"> A) yuzdagi oddiy temiratki 	<ol style="list-style-type: none"> 1. Cellulit A) utkir streptoqoqli 2. Vulgar ektima
III. Strepto-stafilodermiya	
<ol style="list-style-type: none"> 1. Impetigo oddiy (aralash) 	<ol style="list-style-type: none"> 1. Susunkali kechuvchi yarali piodermiya 2. Shankriform piodermiya

Follikulit – soch follikulasini yiringli yalliglanishi
Ajratishadi ostiofollikulit, yuza va chuqur follikulit

Oddiy sikoz ko'pincha erkaklarda uchraydigan kasallik
Bu jarayenda muylov , soqol va boshqa zonadagi
sochlar shikastlanadi



Furunkul – chuqur stafilodermiya. Bu kasallikda soch follikulasini yiringli nekrotik shikastlanish kuzatiladi. Jarayon granulaciya to'qimalari hisobiga chegaralanadi



Karbunkul -bir nechta soch follikullarning yiringli nekrotik yalliglanishi . Yalliglanuvchi infiltrat paydo bo'ladi, derma , gipodermada joylashadi,kattaligi bolalar kaftiday bo'lishi mumkin Bir nechta yiringli nekrotik yalliglanuvchi tugunlar bo'ladi



Gidradenit apokrin bezlarni yiringli yalliglanishi.
Stafilokokklar bezlarning yuliga teridagi travmalar orqali
(kupincha soch olinganda kiradin.



Chaqaloqlarning epidemik porsildoq yarasi Kuponcha bola tugilganda 3-5 kundan keyin paydo buladi - kontagioz yuza stafilodermiya

Impetigo – kontagioz kasallik . Nofolikular , qopqogi ilvillagan , atrofida qizil gardishi ,ishida seroz yiringli suyuqligi bor Pufaklar paydo buladi



Ektima – terini chuqur streptokokkli shiqastlanishi. Bu chuqur yara chetlari kutarilib turadi, tubi yumshoq yiringli , atrofida ogriqli yalliglanuvchi infiltrat kuzatiladi

Surunkali kechuvchi **vegetatsiyali** piodermiya - chuqur piodermiyaning surunkali kechuvchi aralash strepto-stafilokokkli turi .

Shankriform piodermiya – surunkali kechuvchi yarali piodermiya . Zaxmdagi shankrni eslatadi



**E`TIBORINGIZ UCHUN
RAXMAT**