

The image features a white background with a large white circle on the left side. A gold triangle is positioned on the right side, with its hypotenuse facing the circle. The text 'Acute respiratory diseases in children' is centered in the white space between the circle and the triangle.

Acute respiratory diseases in children

Plan of the lecture

1.

2.

3.

4.


5.

6.

7.

RD

ARD is etiologically heterogeneous group of infectious diseases with similar epidemiologic and clinic characteristics.



- **Acute respiratory infection** (ARI) is a common clinical condition of lower respiratory tract infection (LRTI) in children.
- **ARI** is the leading cause of hospitalization in the first 3 y.o.
- **ARI** is a major cause of morbidity in the industry.
- **ARI** is a major cause of mortality in the industry.

Etiology of ARD



ia,

As a rule ARD course isn't severe and rarely produce complications, but sometimes it can initialize another pathologies.

Among respiratory viral diseases the most severe course is in influenza or adenoviral infections, RS viruses or parainfluenza type 3. It's quite commonly accompanied by bacterial infection that worsen condition and prognosis for life.

Transmission mechanism in ARD



• **Transmission mechanism**
• **Transmission routes**
• **Transmission dynamics**
• **Transmission control**




• **Transmission routes**
• **Transmission dynamics**
• **Transmission control**

Susceptibility for ARD infection is universal, but is more prominent in age of 6 mo to 3 y.o. It can be explained by absence of previous contact with these microorganisms and absence of active immunity. Growing children get this immunity and lower their morbidity.

Postinfective specific immunity has its own peculiarities depending on etiology of disease. Influenza or vaccination develop lifelong immunity but viral drift (i.e. not significant antigen changes) raise susceptibility of population and seasonal morbidity sometimes even epidemic. Influenza virus A except drift capable for spontaneous mutations and recombination of RNA fragments (so called antigen shift). Due to this pandemic can appear periodically (once per 10-40 years), when all world population can be affected by these pathogenes.

The total viral serotypes count is about 180 and they cause respiratory tract affection in 95 %



50%
infection
surface

20%
conjugate


15%
antitopic

10%
of

5%
first

its

All viruses produce very similar clinical picture – catarrhal events, running nose, cough and hyperthermia. But some peculiarities exist in various viruses diseases. For instance: **adenoviruses** can cause tonsillitis (frequently with thin coating on tonsils), produce lymphadenopathy, prolonged course of intoxication and fever. **Enteroviruses** can produce herpangina. **Parainfluenza** viruses are the most frequent reason of laryngitis and stridor in children. **RS** viruses produce obstructive bronchitis or bronchiolitis in infants.



Except mechanical defending mechanism, respiratory tract is protected by immune system.

-
-
-
-
-



Neonates after birth are defended by adequate immune response. Besides this they are protected by mother's Ig for 3 mo. But infants has peculiarities of immune system.

Polynuclear neutrophils are able to perform phagocytosis but their mobilization is 2-3 times lower than in adults.

Cytotoxic activity of NK is significantly lower than in adults.

Production of IgM is the same as in adults but secretion of IgA and IgG and reach the proper level is only at 5-7 years old.

Interferon secretion is 10 times less than in adults. Deficiency of IL-2 predispose to Th-2 type of answer and efficient Th-1 way of defending as Th-2 induce secretion of IgE and predispose to atopy.

Fever is the protective- accommodate reaction of organism caused by pathologic agents and characterized by remodeling of thermoregulation process with elevation of body T and stimulation of natural organism reactivity


-
-
-
-

Indications for antipyretic medications

10/10

A decorative graphic consisting of a dashed line that starts from the top left and extends towards the bottom right. To the right of this line, there is a large, curved, gold-colored shape that resembles a stylized arrow or a decorative element.

Risk group for complications due to fever



Hyperthermic syndrome is a pathologic type of fever when a fast raising of body T is accompanied with microvasculature metabolic impairment and progressive dysfunction of essential organs

Main signs of hyperthermia condition:



• Increase in body temperature
• Lasts for 6 hours



• Irritation

Medication choice in fever are



on


nd 2)

e (

Paracetamol is the most safe antipyretic drug. It's dosage is ***10-15 mg/kg tid or 4 times /day.*** Daily dosage mustn't exceed 60mg/kg. Sirup forms of paracetamol start its effect after 30-60 min after admission; In suppositories – effect is realized 2-3 hours later. They are convenient for night time.

Ibuprofen dosage is ***5-10 mg/kg tid.***

Lytic mixture is prescribed only for hyperthermia condition and “pale” fever IM



If child has generalized convulsions it's necessary




any br

can

11

To relief convulsions prescribe parenterally



Toxic syndrome –(acute infectious toxicosis, neurotoxicosis, toxic encephalopathy) is typical for initial period and has several phases.

Transforming of one phase into another can be seen if child don't get proper treatment.

Initial phase Child is apathic, refuse feeding, don't smile, sometimes is irritated, pale with bluish discoloration under the eyes. His sleeping is disturbed, regurgitation or even vomiting can appear. Tachycardia isn't correlated with T, muscle dystonia, contractility of muscle groups, not stable nystagmus can be find.


Irritative phase Nocturnal agitation, painful crying, fast raising of T, tachypnoe and tachycardia, elevation of BP are common signs of second stage Neurologic symptoms appear like tremor and seizures, meningism symptoms.

Hypotonic phase Irritation subsides by adynamia sopor, decreasing of BP muffled heart sound, depressed respiration, tonic convulsions with apnoe.

Deep coma phase Child is slightly react or don't to pain, T decreased. Respiration become aperiodic, hasping type respiration, bradycardia. Skin becomes grayish with marmoreal discoloration due to vascular picture, hemorrhagic rash can appear, DIC syndrome can produce bleeding. Child can die without proper emergency aid.

Typical for toxicosis changes (edema, stasis, hemorrhages, acute dystrophy and alteration) will more visible in systems and organs impaired beforehand. Dominating syndrome like encephalopathic, cardiac hemorrhagic, kidney failure, respiratory distress syndrome will be developed in locus minoris. Such conditions as lost of conscience, prolonged convulsions, signs of brain hypoxia, cardiac failure, hemorrhagic syndrome, kidney failure need emergency treatment.

Toxicosis treatment



Neuro-vegetative protection is performed taking into account such rules:

- **Prophylactic** - aimed at preventing the development of neuro-vegetative disorders.
- **Protonotropic** - aimed at restoring the normal function of the proton pump.
- **Neurotropic** - aimed at restoring the normal function of the neuron.
- **Neuroprotective** - aimed at protecting the neuron from damage.


Typical symptoms of stridor




1)

2)

Stridor degrees



Treatment of stridor (only in hospital!)



in
ages
phala

or

Indications for invasive treatment

-
-
-

of
ent

Clinical peculiarities and signs of ARD

- Allergic rhinitis is characterized by a seasonal or perennial course, with a clear relationship to allergens (pollen, dust, mold, etc.).
- The main clinical signs are: sneezing, nasal itching, watery discharge, and nasal congestion.
- The discharge is typically clear and watery, and is often associated with itchy eyes, nose, and throat.
- The symptoms are usually relieved by antihistamines and nasal corticosteroids.
- The condition is often associated with asthma and allergic conjunctivitis.
- The diagnosis is based on the clinical history and physical examination, and is confirmed by skin prick tests or specific IgE tests.
- The treatment is aimed at avoiding allergens and using medications to relieve symptoms.
- The prognosis is generally good, with symptoms often resolving with appropriate treatment.

Rhinitis treatment is symptomatic:



150

r

tion

for

Pharyngitis - mucous layer inflammation of pharynx. It is frequently combined with rhinitis and is called nasopharyngitis – the most frequent syndrome in ARD.


Symptoms: sudden tickling feeling in the throat dryness, sore throat while swallowing or taking meals. Common condition is usually normal or slightly impaired, body T can be elevated or not. Prognosis is good. Recovery usually in 5-7 days.

Pharyngitis treatment



er


Etiotropic therapy in ARD




Research shows that etiotropic therapy is more effective when initiated before the start of treatment with antihistamines.

- The use of etiotropic therapy is increasing in patients with allergic rhinitis.
- The use of etiotropic therapy is increasing in patients with allergic rhinitis.
- The use of etiotropic therapy is increasing in patients with allergic rhinitis.


Etiotropic therapy in ARD



Etiotropic therapy in ARD




Etiotropic therapy in ARD




50% unknown etiology. The etiology of ARD is unknown in 50% of cases. The etiology of ARD is viral in 25% of cases. The etiology of ARD is bacterial in 15% of cases. The etiology of ARD is fungal in 10% of cases.

Indications for antibiotics in ARD



Control questions



of late

men of

n.

itis.

ical