

# Staphylococcus

### Classification

• Family Micrococcaceae

• Genus Micrococcus and Staphylococcus

Species

more than 20 species S. aureus

S. saprophyticus

S. epidermidis

M. luteus

### **Gram-Positive Cocci**

FAMILY Streptococcaceae (catalase negative)

Group A: β-hemolytic Streptococcus pyogenes

Group B: β-hemolytic (occasionally α or γ) S. agalactiae

Group C: β-hemolytic (α or γ) S. anginosus, S. equismilis

Group D: α or γ hemolytic (β) S. bovis

Group F: β-hemolytic S. anginosus

Group G: β -hemolytic S. anginosus

Viridans streptococci: (no group specific CHO)

α or γ hemolytic S. mutans and

S. salivarius, S. sanguis, S. mitis and S. milleri groups

Streptococcus pneumoniae (no group CHO)(α-hemolytic)

Enterococcus (Group D CHO) γ hemolytic (α or β)


Enterococcus faecalis, E. faecium

FAMILY Micrococcaceae (catalase positive)

Coagulase-positive Staphylococcus aureus


Coag.-neg. Staphylococcus epidermidis, S. saprophyticus

Morphology




Morphology


Staph vs. Strep




Gram-positive cocci in clusters













TABLE 22-2. Staphylococcus, Micrococcus, Stomatococcus, and Alloiococcus and Their Diseases

Organism	Diseases
Staphylococcus aureus	Toxin-mediated (food poisoning, toxic shock syndrome); cutaneous (impetigo, folliculitis, furuncles, carbuncles, wound infections); other (bacteremia, endocarditis, pneumonia, empyema, osteomyelitis, septic arthritis)
Staphylococcus epidermidis	Bacteremia; endocarditis; surgical wounds; urinary tract infections; opportunistic infections of catheters, shunts, prosthetic devices, and peritoneal dialysates
Staphylococcus saprophyticus	Urinary tract infections, opportunistic infections
Staphylococcus capitis	Bacteremia, endocarditis, urinary tract infections, wound infections, pneumonia, bone and joint infections, opportunistic infections
Staphylococcus haemolyticus	Bacteremia, endocarditis, urinary tract infections, wound infections, and opportunistic infections
Micrococcus spp.	Opportunistic infections
Stomatococcus mucilaginosus	Bacteremia, endocarditis, opportunistic infections
Alloiococcus otitidis	Chronic middle ear infections



## See Overheads

~~~~~~~

TSS
Foodborne Intoxication

~~~~~~~

### TABLE 22-3. Staphylococcus aureus Virulence Factors

A ZONENEUE BEZEREN	
Virulence Factors	Biologic Effects
Structural Components	
Capsule	Inhibits chemotaxis and phagocytosis; inhibits proliferation of mononuclear cells; facili- tates adherence to foreign bodies
Peptidoglycan	Provides osmotic stability; stimulates production of endogenous pyrogen (endotoxin-like activity); leukocyte chemoattractant (abscess formation); inhibits phagocytosis
Teichoic acid	Regulates cationic concentration at cell membrane; binds to fibronectin
Protein A	Inhibits antibody-mediated clearance by binding IgG <sub>1</sub> , IgG <sub>2</sub> , and IgG <sub>4</sub> Fc receptors; leukocyte chemoattractant; anticomplementary
Cytoplasmic membrane	Osmotic barrier; regulates transport into and out of cell; site of biosynthetic and respiratory enzymes
Toxins	
Cytotoxins ( $\alpha$ , $\beta$ , $\delta$ , $\gamma$ , P-V leukocidin)	Toxic for many cells, including leukocytes, erythrocytes, macrophages, platelets, and fibroblasts
Exfoliative toxins (ETA, ETB)	Serine proteases that split the intercellular bridges in the stratum granulosum epidermis
Enterotoxins (A–E, G–I)	Superantigens (stimulates proliferation of T cells and release of cytokines); stimulates release of inflammatory mediators in mast cells, increasing intestinal peristalsis and fluid loss, as well as nausea and vomiting
Toxic Shock Syndrome Toxin-1	Superantigen (stimulates proliferation of T cells and release of cytokines); produces leakage or cellular destruction of endothelial cells
Enzymes	and the second of the second o
Coagulase	Converts fibrinogen to fibrin
Catalase	Catalyzes removal of hydrogen peroxide
Hyaluronidase	Hydrolyzes hyaluronic acids in connective tissue, promoting the spread of staphylococci in tissue
Fibrinolysin	Dissolves fibrin clots
Lipases	Hydrolyzes lipids
Nucleases	Hydrolyzes DNA
Penicillinase	Hydrolyzes penicillins

### **MICROBIAL PATHOGEN**

**ADHESIN** 

**RECEPTOR** 

Staphylococcus aureus

Staphylococcus spp.

Group A Streptococcus

Streptococcus pneumoniae

Escherichia coli

Other Enterobacteriaceae

Neisseria gonorrhoeae

Treponema pallidum

Chlamydia spp.

Mycoplasma pneumoniae

Vibrio cholerae

Lipoteichoic acid

Slime layer

LTA-M protein complex Fibronectin

Protein

Type 1 fimbriae

CFA 1 fimbriae

P fimbriae

Type 1 fimbriae

**Fimbriae** 

 $P_1, P_2, P_3$ 

Cell surface lectin

Protein P1

Type 4 pili

Unknown

Unknown

**D-Mannose** 

GM ganglioside

P blood grp glycolipid

N-acetylhexosamine-gal

**D-Mannose** 

GD<sub>1</sub> ganglioside

Fibronectin

N-acetylglucosamine

Sialic acid

Fucose and mannose

### Cell-Associated Virulence Factors

- Capsule or slime layer (glycocalyx)
- Peptidoglycan (PG)
- Teichoic acid is covalently linked to PG and is species specific:
  - S. aureus ribitol teichoic acid (polysaccharide A)
  - S. epidermidis glycerol teichoic acid (polysaccharide B)
- Protein A is covalently linked to PG
- Clumping factor (bound coagulase)

# Virulence Factors Extracellular Enzymes

- Coagulases (bound or free)
  - ✓ Antigenic
- Hyaluronidase
  - ✓ "spreading factor" of S. aureus
- Nuclease
  - Cleaves DNA and RNA in S. aureus
- Protease
  - ✓ Staphylokinase (fibrinolysin)
- Lipases
- Esterases

## Virulence Factors: Exotoxins

- Cytolytic (cytotoxins; cytolysins)
  - ✓ Alpha toxin hemolysin
    - Reacts with RBCs
  - ✓ Beta toxin
    - Sphingomyelinase
  - ✓ Gamma toxin
    - Hemolytic activity
  - ✓ Delta toxin
    - Cytopathic for:
      - RBCs
      - Macrophages
      - Lymphocytes
      - Neutrophils
      - Platelets
    - Enterotoxic activity
  - ✓ Leukocidin

# Virulence Factors: Exotoxins

- Enterotoxin
- Exfoliative toxin (epidermolytic toxin)
- Pyrogenic exotoxins

# **Pathogenesis**

- Pass skin first line of defense
  - Benign infection
    - Phagocytosis
    - Antibody
    - Inflammatory response
  - Chronic infections
    - Delayed hypersensitivity

### Clinical Manifestations/Disease

### SKIN

- ✓ folliculitis
- ✓ boils (furuncles)
- ✓ carbuncles
- ✓ impetigo (bullous & pustular)
- scalded skin syndrome
  - Neonates and children under 4 years







### Clinical Manifestations/Disease

- Other infections
  - Primary staphylococcal pneumonia
  - ✔ Food poisoning vs. foodborne disease
  - ✓ Toxic shock syndrome



# Metastatic Infections

- Bacteremia
- Osteomyelitis
  - ✓ disease of growing bone
  - Pulmonary and cardiovascular infection





# Coagulase-Negative Staphylococci

- Staphylococcus epidermidis
  - S. saprophyticus

### **BOX 22-2.** Summary of Coagulase-Negative Staphylococcal Infections

#### Physiology and Structure

Gram-positive cocci arranged in clusters.

Facultative anaerobe (capable of aerobic and anaerobic growth).

Catalase-positive but coagulase-negative.

Species-specific teichoic acid.

Capsule ("slime" layer) present.

#### Virulence

Refer to Table 22-3.

### **Epidemiology**

Normal human flora on skin and mucosal surfaces.

Organisms can survive on dry surfaces for long periods.

Person-to-person spread through direct contact or exposure to contaminated fomites (although most infections are with the patient's own organisms).

Patients at risk are those with foreign bodies (e.g., suture, prosthesis, shunt, catheter).

The organisms are ubiquitous, so there are no geographic or seasonal limitations.

#### Diseases

Diseases—see Table 22-2.

Catheter-related bacteremia.

Subacute endocarditis associated with previously damaged or artificial heart valve.

Central nervous system shunt infection.

Surgical wound infection when a foreign body (e.g., suture, prosthesis, medical hardware) is present.

#### Diagnosis

As with S. aureus infections.

#### Treatment, Control, and Prevention

The antibiotics of choice are oxacillin (or other penicillinaseresistant penicillin) or vancomycin for oxacillin-resistant strains.

Removal of the foreign body is frequently required for successful treatment.


Prompt treatment for endocarditis or shunt infections is necessary to prevent further tissue damage or immune complex formation.

Maintenance of sterile intravascular catheters helps prevent infections.



### Staphylococcal Lab ID & Diagnostic Tests

- Microscopic
- Lab isolation
- Coagulase positive
  - ✓ S. aureus











### **Differential Characteristics**

### Catalase

 $2H_2O_2 \square O_2 + 2H_2O$ Streptococci vs. Staphylococci










# Differential Characteristics


Coagulase

Fibrinogen 

Fibrin







A positive reaction is indicated by coagulation of the plasma.

Coagulase POS

If the cells did not produce coagulase, the plasma remains fluid.

Coagulase NEG

### **Treatment**

- Drain infected area
- Deep/metastatic infections
  - ✓ semi-synthetic penicllins
  - ✓ cephalosporins
  - ✓ erythromycin
  - ✓ clindamycin
- Endocarditis
  - ✓ semi-synthetic penicillin + an aminoglycoside

### Prevention

- Carrier status prevents complete control
- Proper hygiene, segregation of carrier from highly susceptible individuals
- Good aseptic techniques when handling surgical instruments
- Control of nosocomial infections





### **Gram-Positive Cocci**

FAMILY Streptococcaceae (catalase negative)

Group A: β-hemolytic Streptococcus pyogenes

Group B: β-hemolytic (occasionally α or γ) S. agalactiae

Group C: β-hemolytic (α or γ) S. anginosus, S. equismilis

Group D: α or γ hemolytic (β) S. bovis

Group F: β-hemolytic S. anginosus

Group G: β -hemolytic S. anginosus

Viridans streptococci: (no group specific CHO)

α or γ hemolytic S. mutans and

S. salivarius, S. sanguis, S. mitis and S. milleri groups

Streptococcus pneumoniae (no group CHO)(α-hemolytic)

Enterococcus (Group D CHO) γ hemolytic (α or β)

Enterococcus faecalis, E. faecium

FAMILY Micrococcaceae (catalase positive)
Coagulase-positive Staphylococcus aureus
Coag.-neg. Staphylococcus epidermidis, S. saprophyticus

REVIEW

### Which features are only found in S. aureus?




TABLE 22-2. Staphylococcus, Micrococcus, Stomatococcus, and Alloiococcus and Their Diseases

Organism	Diseases
Staphylococcus aureus	Toxin-mediated (food poisoning, toxic shock syndrome); cutaneous (impetigo, folliculitis, furuncles, carbuncles, wound infections); other (bacteremia, endocarditis, pneumonia, empyema, osteomyelitis, septic arthritis)
Staphylococcus epidermidis	Bacteremia; endocarditis; surgical wounds; urinary tract infections; opportunistic infections of catheters, shunts, prosthetic devices, and peritoneal dialysates
Staphylococcus saprophyticus	Urinary tract infections, opportunistic infections
Staphylococcus capitis	Bacteremia, endocarditis, urinary tract infections, wound infections, pneumonia, bone and joint infections, opportunistic infections
Staphylococcus haemolyticus	Bacteremia, endocarditis, urinary tract infections, wound infections, and opportunistic infections
Micrococcus spp.	Opportunistic infections
Stomatococcus mucilaginosus	Bacteremia, endocarditis, opportunistic infections
Alloiococcus otitidis	Chronic middle ear infections



#### Physiology and Structure

Gram-positive cocci aranged in clusters.

Facultative anaerobe (capable of aerobic and anaerobic growth).

Coagulase ("clumping factor") positive (key identification test).

Species-specific teichoic acid.

Protein A (useful for identification).

Capsule.

#### Virulence

Virulence factors—see Table 22-3.

#### **Epidemiology**

Normal flora on human skin and mucosal surfaces.

Organisms can survive on dry surfaces for long periods (owing to thickened peptidoglycan layer and absence of outer membrane—characteristics of all gram-positive bacteria).

Person-to-person spread through direct contact or exposure to contaminated fomites (e.g., bed linens, clothing).

Risk factors include presence of a foreign body (e.g., splinter, suture, prosthesis, catheter), previous surgical procedure, use of antibiotics that suppress the normal microbial flora).

Patients at risk for specific diseases include menstruating women (toxic shock syndrome), infants (scalded skin syndrome), young children with poor personal hygiene (impetigo and other cutaneous infections), intravascular catheters (bacteremia and endocarditis); patients with compromised pulmonary function or an antecedent viral respiratory infection (pneumonia).

Infections are found worldwide and generally with no seasonal prevalence (except that food poisoning is more common in summer and during late-year holidays).

#### Diseases

Refer to Table 22-2.

Toxin-mediated diseases include food poisoning, toxic shock syndrome, and scalded skin syndrome.

Pyogenic diseases include impetigo, folliculitis, furuncles, carbuncles, and wound infections.

Other systemic diseases (frequently associated with bacteremia) include pneumonia (typically after viral respiratory infection), empyema (complication of pneumonia or surgical intervention), septic arthritis, osteomyelitis, acute endocarditis, and catheter-related bacteremia.

#### Diagnosis

Microscopy is useful for pyogenic infections but not blood stream infections or toxin-mediated infections.

Staphylococci grow rapidly when cultured on nonselective media.

Detection of staphylococcal antigens by serology are generally of low value.

#### Treatment, Control, and Prevention

The antibiotics of choice are oxacillin (or other penicillinase-resistant penicillin) or vancomycin for oxacillin-resistant strains.

The focus of infection (e.g., abscess) must be identified and drained.

Treatment is symptomatic for patients with food poisoning (although the source of infection should be identified so that appropriate preventive procedures can be enacted).

Proper cleansing of wounds and use of disinfectant help prevent infections.

Thorough hand washing and covering of exposed skin helps medical personnel prevent infection or spread to other patients.











**REVIEW** 

### TABLE 22-3. Staphylococcus aureus Virulence Factors

Virulence Factors	Biologic Effects
Structural Components	
Capsule	Inhibits chemotaxis and phagocytosis; inhibits proliferation of mononuclear cells; facili- tates adherence to foreign bodies
Peptidoglycan	Provides osmotic stability; stimulates production of endogenous pyrogen (endotoxin-like activity); leukocyte chemoattractant (abscess formation); inhibits phagocytosis
Teichoic acid	Regulates cationic concentration at cell membrane; binds to fibronectin
Protein A	Inhibits antibody-mediated clearance by binding IgG <sub>1</sub> , IgG <sub>2</sub> , and IgG <sub>4</sub> Fc receptors; leukocyte chemoattractant; anticomplementary
Cytoplasmic membrane	Osmotic barrier; regulates transport into and out of cell; site of biosynthetic and respiratory enzymes
Toxins	
Cytotoxins ( $\alpha$ , $\beta$ , $\delta$ , $\gamma$ , P-V leukocidin)	Toxic for many cells, including leukocytes, erythrocytes, macrophages, platelets, and fibroblasts
Exfoliative toxins (ETA, ETB)	Serine proteases that split the intercellular bridges in the stratum granulosum epidermis
Enterotoxins (A-E, G-I)	Superantigens (stimulates proliferation of T cells and release of cytokines); stimulates release of inflammatory mediators in mast cells, increasing intestinal peristalsis and fluid loss, as well as nausea and vomiting
Toxic Shock Syndrome Toxin-1	Superantigen (stimulates proliferation of T cells and release of cytokines); produces leakage or cellular destruction of endothelial cells
Enzymes	
Coagulase	Converts fibrinogen to fibrin
Catalase	Catalyzes removal of hydrogen peroxide
Hyaluronidase	Hydrolyzes hyaluronic acids in connective tissue, promoting the spread of staphylococci in tissue
Fibrinolysin	Dissolves fibrin clots
Lipases	Hydrolyzes lipids
Nucleases	Hydrolyzes DNA
Penicillinase	Hydrolyzes penicillins REVIE

### **BOX 22-2.** Summary of Coagulase-Negative Staphylococcal Infections

#### Physiology and Structure

Gram-positive cocci arranged in clusters.

Facultative anaerobe (capable of aerobic and anaerobic growth).

Catalase-positive but coagulase-negative.

Species-specific teichoic acid.

Capsule ("slime" layer) present.

#### Virulence

Refer to Table 22-3.

### **Epidemiology**

Normal human flora on skin and mucosal surfaces.

Organisms can survive on dry surfaces for long periods.

Person-to-person spread through direct contact or exposure to contaminated fomites (although most infections are with the patient's own organisms).

Patients at risk are those with foreign bodies (e.g., suture, prosthesis, shunt, catheter).

The organisms are ubiquitous, so there are no geographic or seasonal limitations.

#### Diseases

Diseases—see Table 22-2.

Catheter-related bacteremia.

Subacute endocarditis associated with previously damaged or artificial heart valve.

Central nervous system shunt infection.

Surgical wound infection when a foreign body (e.g., suture, prosthesis, medical hardware) is present.

#### Diagnosis

As with S. aureus infections.

#### Treatment, Control, and Prevention

The antibiotics of choice are oxacillin (or other penicillinaseresistant penicillin) or vancomycin for oxacillin-resistant strains.

Removal of the foreign body is frequently required for successful treatment.

Prompt treatment for endocarditis or shunt infections is necessary to prevent further tissue damage or immune complex formation.

Maintenance of sterile intravascular catheters helps prevent infections.

