


Physiology of Bacteria

Department of Microbiology, Virology
and Immunology



Microbial Metabolism

- The primary function of all living cells is to grow and reproduce
- Growth & reproduction rely on the outcome of chemical reactions in the cells
- The sum of all cellular chemical reactions is referred to as **metabolism**



Microbial Metabolism

- The metabolic process that involves the **degradation** of chemical components is called **catabolism**
- The synthesis of chemical components is called **anabolism** or **biosynthesis**



- Most metabolic processes in the cell would take forever if it were not for **enzymes**.
- **Enzymes** are proteins that have molecular weights ranging from 600 to 12 000.
 - Their function is to speed up the various chemical reactions that occur in the cell.
 - Molecules that speed up chemical reactions are called **catalysts**.
 - Enzymes often cannot function alone and require additional molecules, called **cofactors**, to enhance activity



Classification of enzymes

- **Oxidoreductases** are involved in electron (hydrogen) transfer reactions.
- **Transferases** transfer specific groups such as aldehydes or phosphates from one substrate to another.
- **Hydrolyses** add water across chemical bonds to be cleaved or hydrolyzed.



Classification of enzymes

- **Lyases** remove chemical groups from substrates, forming double bonds, or add chemical groups to double bonds.
- **Isomerases** rearrange certain compounds to produce molecules having the same groups of atoms, but in different arrangements.
- **Ligases** produce bonds accompanied by the cleavage of ATP.



Classification of enzymes

- Enzymes synthesized by the cell remain within the cell to carry out specific reactions and are called **endoenzymes**
- Enzymes released from the cell into the surrounding environment and are called **exoenzymes**



Classification of enzymes

- **Pathogenicity enzymes** - are enzymes that damage cells and tissues.
 - **Coagulase** –enables the organisms to clot plasma to form a sticky coat of fibrin around themselves for protection from phagocytes and other body defense mechanisms (Staphylococcus).
 - **Kinases** –referred to as fibrinolysin, kinase has opposite effect of coagulase. Streptokinase, for example, lyses fibrin clots, thus enabling streptococci to invade and spread throughout the body.



Classification of enzymes

- **Hyaluronidase** –enables pathogens to spread through connective tissue by breaking down hyaluronic acid, the “cement” that holds tissue cells together (Staphylococcus, Streptococcus and Clostridium).
- **Collagenase**- This enzyme breaks down collagen, the supportive protein founding tendons, cartilage and bones. Cl. perfringens a major cause of gas gangrene, spreads deeply within the body by secreting both collagenase and hyaluronidase.



Classification of enzymes


- **Hemolysin-** enzyme that cause damage to the host's red blood cells. In the laboratory, hemolysis of the red blood cells in the blood agar is useful for identifying types of *Staphylococcus* and *Streptococcus*.
- **Lecithinase** – one of the toxins produced by *Staphylococcus aureus*, which breaks down phospholipids collectively referred to as lecithin.
- **Leukocidin-** enzyme secreted some *Staphylococcus aureus* causes destruction



Growth & Multiplication of Bacteria

- Bacteria divide by binary fission
- Bacterial cell divides to form two daughter cells
- Nuclear division precedes cell division & in a growing population many cells carrying two nuclear bodies can be seen





- The interval of time between two cell division, or the time required for a bacterium to give rise to two daughter cells under optimum conditions, is known as the generation time or population doubling time





- In many medically important bacteria, the generation time is about 20 minutes
- Some bacteria are slow-growing
 - Tubercle bacilli the generation time is about 20 hours
 - Lepra bacilli about 20 days
- Bacteria reproduce so rapidly & by geometric progression, a single bacterial cell can theoretically give rise to 10^{21} progeny in 24 hours, with a mass of approximately 4,000 tones!



- When bacteria are grown in a vessel of liquid medium (batch culture), multiplication is arrested after a few cell divisions due to depletion of nutrients or accumulation of toxic products
- When pathogenic bacteria multiply in host tissues, the situation may be intermediate between a batch culture & a continuous culture
- Bacteria growing on solid media form colonies
- Each colony represents a clone of cells derived from a single parent cell
- In liquid media, growth is diffuse




Bacterial cell Growth Curve

- **A- Lag phase**

- Immediately following the seeding of a culture medium
- This initial period is the time required for adaptation to the new environment
- There is no increase in numbers, though there may be an increase in the size of the cells

- **B- Log (logarithmic) or exponential phase**

- The cells start dividing & their numbers increase exponentially or by geometric progression




Bacterial cell Growth Curve

- **C- Stationary phase**
 - After a period of exponential growth, cell division stops due to depletion of nutrients & accumulation of toxic products
 - The viable count remains stationary as an equilibrium exists between the dying cells and the newly formed cells
- **D- Phase of Decline**
 - Population decreases due to cell death



Bacterial cell Growth Curve



Nutritional requirements

- Microorganisms also depend on an available source of chemical nutrients. Microorganisms are often grouped according to their energy source and their source of carbon.
- a. **Energy source**
 - 1. **Phototrophs** use radiant energy (light) as their primary energy source.
 - 2. **Chemotrophs** use the oxidation and reduction of chemical compounds as their primary energy source.
- b. **Carbon source**
- Based on their source of carbon bacteria can be classified as autotrophs or heterotrophs.
 - 1. **Autotrophs**: require only carbon dioxide as a carbon source. An autotroph can synthesize organic molecules from inorganic nutrients.
 - 2. **Heterotrophs**: require organic forms of carbon. A Heterotroph cannot synthesize organic molecules from inorganic nutrients.



- All organisms in nature can be placed into one of four separate groups: photoautotrophs, photoheterotrophs, chemoautotrophs, and chemoheterotrophs.
- 1. **Photoautotrophs** use **light** as an energy source and **carbon dioxide** as their main carbon source. They include photosynthetic bacteria (green sulfur bacteria, purple sulfur bacteria, and cyanobacteria), algae, and green plants. Photoautotrophs transform carbon dioxide and water into carbohydrates and oxygen gas through **photosynthesis**.
- 2. **Photoheterotrophs** use **light** as an energy source but cannot convert carbon dioxide into energy.. They include the green nonsulfur bacteria and the purple nonsulfur bacteria.
- 3. **Chemolithoautotrophs** use **inorganic compounds** such as hydrogen sulfide, sulfur, ammonia, nitrites, hydrogen gas, or iron as an energy source and **carbon dioxide** as their main carbon source.
- 4. **Chemoorganoheterotrophs** use **organic compounds** as both an energy source and a carbon source. **Saprophytes** live on dead organic matter while **parasites** get their nutrients from a living host. Most bacteria, & all protozoans, fungi, and animals are chemoorganoheterotrophs.



Nutritional requirements

- d. **Minerals**
- 1. **sulfur**-Sulfur is needed to synthesize sulfur-containing amino acids and certain vitamins.
- 2. **phosphorus** -Phosphorus is needed to synthesize phospholipids (*def*), DNA, RNA, and ATP (*def*). Phosphate ions are the primary source of phosphorus.
- 3. **potassium, magnesium, and calcium**-These are required for certain enzymes to function as well as additional functions.
- 4. **iron**-Iron is a part of certain enzymes.
- 5. **trace elements** -Trace elements are elements required in very minute amounts, and like potassium, magnesium, calcium, and iron, they usually function as **cofactors** (*def*) in enzyme reactions. They include sodium, zinc, copper, molybdenum, manganese, and cobalt ions. Cofactors usually function as electron donors or electron acceptors during enzyme reactions.



Nutritional requirements

- e. **Water**

- f. **Growth factors**

Growth factors are organic compounds such as amino acids (*def*), purines (*def*), pyrimidines (*def*), and vitamins (*def*) that a cell must have for growth but cannot synthesize itself. Organisms having complex nutritional requirements and needing many growth factors are said to be **fastidious**.



Oxygen Requirements

- Depending on the influence of oxygen on growth and viability, bacteria are divided into **aerobes & anaerobes**
- Aerobic bacteria require oxygen for growth

Aerobic bacteria



obligate aerobes

(*Vibrio cholerae*)



Oxygen Requirements

- Anaerobic bacteria grow only in absence of oxygen

Anaerobic bacteria

obligate anaerobe

(clostridia)

facultative anaerobes

(most of medically
important bacteria)



Oxygen Requirements

- **Obligate anaerobes** are organisms that grow **only** in the absence of oxygen and, in fact, are often inhibited or killed by its presence. They obtain their energy through anaerobic respiration or fermentation.
- **Facultative anaerobes** are organisms that grow with or without oxygen, but generally better with oxygen. They obtain their energy through aerobic respiration if oxygen is present, but use fermentation or anaerobic respiration if it is absent. Most bacteria are facultative anaerobes.



Oxygen Requirements

- **Microaerophiles** are organisms that require a low concentration of oxygen (2% to 10%) for growth, but higher concentrations are inhibitory. They obtain their energy through aerobic respiration.
- **Aerotolerant anaerobes** like obligate anaerobes, cannot use oxygen to transform energy but can grow in its presence. They obtain energy only by fermentation and are known as obligate fermenters.



Physical requirements

- **Temperature**
- 1. **Psychrophiles** are cold-loving bacteria. Their optimum growth temperature is between -5C and 15C. They are usually found in the Arctic and Antarctic regions and in streams fed by glaciers.
- 2. **Mesophiles** are bacteria that grow best at moderate temperatures. Their optimum growth temperature is between 25C and 45C. Most bacteria are mesophilic and include common soil bacteria and bacteria that live in and on the body.



Physical requirements

- 3. **Thermophiles** are heat-loving bacteria. Their optimum growth temperature is between 45C and 70C and are commonly found in hot springs and in compost heaps.
- 4. **Hyperthermophiles** are bacteria that grow at very high temperatures. Their optimum growth temperature is between 70C and 110C (Archae are found growing near hydrothermal vents at great depths in the ocean).



- **pH**
- Microorganisms can be placed in one of the following groups based on their optimum pH requirements:
- 1. Neutrophiles grow best at a pH range of 5 to 8.
- 2. Acidophiles grow best at a pH below 5.5.
- 3. Alkaliphiles grow best at a pH above 8.5.



Culture Media

- A **growth medium** or **culture medium** is a substance in which microorganisms is a substance in which microorganisms or cells is a substance in which microorganisms or cells can grow
- There are two major types of growth media: those used for cell culture There are two major types of growth media: those used for cell culture, which use specific cell types derived from plants or animals, and microbiological culture There are two major types of growth media: those used for cell



Types of Growth Media

- The most common growth media for microorganisms are *nutrient broths* (liquid nutrient medium) or *Lysogeny broth* (LB medium). Bacteria grown in liquid cultures often form colloidal (LB medium). Bacteria grown in liquid cultures often form colloidal suspensions.
- Liquid mediums are often mixed with agar Liquid mediums are often mixed with agar and poured into petri dishes Liquid mediums are often mixed with agar and poured into petri dishes to solidify. These




Types of Growth Media

- **Nutrient media**
 - *Undefined media* (also known as *basal* or *complex media*)
 - *Defined media* (also known as chemical defined media)
 - *Differential medium* some sort of indicator, typically a dye, is added, that allows for the differentiation of particular chemical reactions occurring during growth



Types of Growth Media

- **Selective media**
(are used for the growth of only select microorganisms)



Blood-free, charcoal-based selective medium agar (CSM) for isolation of *Campylobacter*

Types of Growth Media


- **Differential media** or *indicator media* distinguish one microorganism type from another growing on the same media (MacConkey's, Nagler's medium)

This type of media uses the biochemical characteristics of a microorganism growing in the presence of specific nutrients or indicators (such as neutral red, phenol red, eosin y, neutral red, phenol red, eosin y, or methylene blue)



Types of Growth Media


Shigella sp., Escherichia sp., and Proteus sp.



MacConkey Agar



Shigella-Salmonella Agar



Bismuth Sulfite Agar



Brilliant Green Agar

Types of Growth Media

- **Enriched media** contain the nutrients required to support the growth of a wide variety of organisms
 - Blood agar is an enriched medium in which nutritionally rich whole blood supplements the basic nutrients.
 - Chocolate agar is enriched with heat-treated blood (40-45°C), which turns brown and gives the medium the color for which it is named.





Blood agar plates are often used to diagnose infection. On the right is a positive Staphylococcus infection; on the left a positive Streptococcus culture.

Types of Growth Media

- **Transport media** used for the temporary storage of specimens being transported to the laboratory for cultivation. Transport media typically contain only buffers and salt (Stuart's medium for gonococci, buffered glycerol saline for enteric bacilli).
- **Indicator media** contain an indicator which changes colour when a bacterium grows in them (Bismuth sulphite media(*S.typhi*), potassium tellurite(*diphtheria bacilli*)).



Types of Growth Media

- **Sugar Media** used for sugar fermentation (Hiss's serum sugars)
 - The sugar media consist of 1% of the sugar in peptone water along with an appropriate indicator
 - Durham's tube is kept inverted in the sugar tube to detect gas production
- **Anaerobic media** are used to grow anaerobic organisms (Robertson's cooked meat medium)

